


BP-1-102

Chemical Properties

CAS No. : 1334493-07-0
 Formula: C₂₉H₂₇F₅N₂O₆S
 Molecular Weight: 626.59
 Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year
 Actual storage temperature shall be subject to the COA.



Biological Description

Description	BP-1-102 is an orally active, effective, and specific STAT3 inhibitor that binds Stat3 (Kd: 504 nM), blocks Stat3-phosphotyrosine (pTyr) peptide interactions and Stat3 activation (4-6.8 μM), and selectively inhibits the migration, survival, growth, and invasion of Stat3-dependent tumor cells. The BP-1-102-mediated inhibition of aberrantly active Stat3 in tumor cells suppresses the expression of [c-Myc, Bcl-xL, Cyclin D1, Survivin, and VEGF].
Targets(IC50)	STAT
In vitro	BP-1-102 binds Stat3 with an affinity (KD) of 504 nM, inhibits Stat3-phospho-tyrosine (pTyr) peptide interactions and Stat3 activation at 4-6.8 μM, and selectively inhibits the growth, survival, migration, and invasion of Stat3-dependent tumor cells. BP-1-102-mediated inhibition of aberrant Stat3 in tumor cells suppresses the expression of c-Myc, Cyclin D1, Bcl-xL, Survivin, VEGF, and Krüppel-like factor 8, a Stat3 target gene promoting breast tumor cell migration and invasion. Treatment of breast cancer cells with BP-1-102 also blocks Stat3-NF-κB cross-talk, the release of several cytokines, and the phosphorylation of focal adhesion kinase and paxillin, while enhancing E-cadherin expression. BP-1-102 inhibits Stat3 DNA-binding activity in vitro with an IC ₅₀ of 6.8±0.8 μM and preferentially inhibits Stat3-Stat3 over other Stat dimer DNA-binding activities. BP-1-102 exerts minimal or no effect on phospho-Shc, Src, Jak-1/2, Erk1/2, or Akt levels [1].
In vivo	Intravenous or oral gavage delivery of BP-1-102 furnishes micromolar or microgram levels in tumor tissues and inhibits growth of human breast and lung tumor xenografts and modulates Stat3 activity, Stat3 target genes, and soluble factors in vivo. BP-1-102 selectively suppresses growth, survival, malignant transformation, migration, and invasion of malignant cells harboring constitutively active stat3. BP-1-102 is detectable at micromolar concentrations in plasma and in micrograms in tumor tissues[1].
Kinase Assay	Kinase targets are tested with biochemical enzymatic kinase assays using the SelectScreen Kinase Profiling Service to determine IC ₅₀ values. The compounds (XMD8-87) are assayed at 10 concentrations (3-fold serial dilutions starting from 1 μM) at an ATP concentration equal to the ATP K _m [1].
Cell Research	Proliferating cells in 6- or 96-well plates are treated once with 0-30 μM BP-1-102 for 24 h or with 10 μM BP-1-102 for up to 96 h. Viable cells are counted by trypan blue exclusion/phase-contrast microscopy or assessed by a CyQUANT Cell Proliferation Kit. (Only for Reference)

Solubility Information

Solubility	Ethanol: < 1 mg/mL (insoluble or slightly soluble), DMSO: 150 mg/mL (239.39 mM),Sonication is recommended. H2O: < 1 mg/mL (insoluble or slightly soluble), (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 3.3 mg/mL (5.27 mM),Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.5959 mL	7.9797 mL	15.9594 mL
5 mM	0.3192 mL	1.5959 mL	3.1919 mL
10 mM	0.1596 mL	0.798 mL	1.5959 mL
50 mM	0.0319 mL	0.1596 mL	0.3192 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

- Zhang X, et al. Proc Natl Acad Sci U S A. 2012, 109(24):9623-8.
- Zhou B, Yan J, Guo L, et al. Hepatoma cell-intrinsic TLR9 activation induces immune escape through PD-L1 upregulation in hepatocellular carcinoma. Theranostics. 2020, 10(14): 6530.
- He Q R, Tang J J, Chen Z F, et al. Natural Product Trienomycin A is a STAT3 pathway inhibitor that exhibits potent in vitro and in vivo efficacy against pancreatic cancer[J]. Authorea Preprints. 2020
- He, Qiu-Rui, et al. The natural product trienomycin A is a STAT3 pathway inhibitor that exhibits potent in vitro and in vivo efficacy against pancreatic cancer. British Journal of Pharmacology. 178.12 (2021): 2496-2515.
- Chen C, Lu M, Lin S, et al. The nuclear gene rpl18 regulates erythroid maturation via JAK2-STAT3 signaling in zebrafish model of Diamond-Blackfan anemia. Cell Death & Disease. 2020, 11(2): 1-11
- Yan J, Zhou B, Guo L, et al. GOLM1 upregulates expression of PD-L1 through EGFR/STAT3 pathway in hepatocellular carcinoma[J]. American journal of cancer research. 2020, 10(11): 3705.
- Chen C, Lu M, Lin S, et al. The nuclear gene rpl18 regulates erythroid maturation via JAK2-STAT3 signaling in zebrafish model of Diamond-Blackfan anemia[J]. Cell Death & Disease. 2020, 11(2): 1-11.
- Yan J, Zhou B, Guo L, et al. GOLM1 upregulates expression of PD-L1 through EGFR/STAT3 pathway in hepatocellular carcinoma. American journal of cancer research. 2020, 10(11): 3705.
- Zhou B, Yan J, Guo L, et al. Hepatoma cell-intrinsic TLR9 activation induces immune escape through PD-L1 upregulation in hepatocellular carcinoma[J]. Theranostics. 2020, 10(14): 6530.
- Strobel T D, Weber M, Heber N, et al.Revisiting the role of endogenous STAT3 in HPV-positive cervical cancer cells. Journal of Medical Virology.2023, 95(11): e29230.
- Zhang P, Wang X, Yang X, et al.Molecular control of PDPN hi macrophage subset induction by ADAP as a host defense in sepsis.JCI insight.2025

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481