


Radotinib

Chemical Properties

CAS No. :	926037-48-1
Formula:	C ₂₇ H ₂₁ F ₃ N ₈ O
Molecular Weight:	530.5
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



Biological Description

Description	Radotinib (IY-5511), and sometimes referred to by its investigational name IY5511, is a drug for the treatment of different types of Y, most notably Philadelphia chromosome-positive (Ph+) chronic myeloid leukemia (CML) with resistance or intolerance of other tyrosine kinase Bcr-Abl inhibitors
Targets(IC50)	Apoptosis,Bcr-Abl,STAT,JAK
In vitro	In vitro, Radotinib binds BCR-ABL1 and reduces phosphorylation of CrkL, a BCR-ABL1 target protein. Radotinib also effectively inhibits the proliferation of common mutant clones of BCR-ABL1, with the exception of T315I. [1] In AML cells, radotinib significantly decreases the cell viability, promotes differentiation, and induces CD11b expression and apoptosis. In NB4, THP-1, and Kasumi-1 cells, radotinib also induces CD11b expression, and decreases the viability. [2]
Kinase Assay	Autophosphorylation of EGFR in cells: For experiments using cells in culture, A431 cells are treated with various concentrations of Pelitinib for 2.75 hours before co-incubation with 100 ng/mL EGF for 0.25 hour. Cells are washed twice with cold phosphate-buffered saline (PBS) before adding to lysis buffer (10 mM Tris, pH 7.5, 5 mM ethylenediamine tetra-acetic acid (EDTA), 150 mM NaCl, 1% Triton X-100, 1% Sodium deoxycholate, 0.1 % SDS, 1 mM PMSF, 10 mg/mL pepstatin A, 10 mg/mL leupeptin, 20 KIU/mL aprotinin, 2 mM sodium orthovanadate, and 100 mM sodium fluoride) for 20 minutes on ice, before immunoprecipitation and SDS-PAGE-immunoblotting. For immunoprecipitation, cultured cells are placed in cold lysis buffer and immediately homogenized on ice with a polytron with several pulses. The homogenate is first centrifuged at 2500 rpm (20 minutes, 4 °C) and then again at 14,000 rpm in a microcentrifuge (10 minutes, 4 °C). Supernatants (1000 µg protein) are incubated for 2 hours at 4 °C with 15 mL of EGFR polyclonal antibody. After 2 hours, 50 µL of protein G plus/protein A agarose beads is added and incubated with constant rotation for 2 hours at 4 °C. After washing with lysis buffer, beads are boiled for 2 minutes in Laemmli sample buffer. Proteins are then resolved by SDS-PAGE, transferred to immobilon membrane and probed overnight with an anti-phosphotyrosine antibody conjugated with horseradish peroxidase (HRP). Membranes are developed using the ECL reagent. Total EGFR protein is determined by stripping membranes and re-probing with receptor-specific antibodies. Quantitation of bands is done by densitometry, using ImageQuant software with a Molecular Dynamics laser transmittance scanner.

A DRUG SCREENING EXPERT

Cell Research	Cells are seeded in 96-well plates at a density of 2×10^4 cells/ml with 100 μ L of medium per well and then incubated with various concentrations of radotinib (0, 1, 10, and 100 μ M) for 72 h at 37°C. The CellTiter 96 solution (20 μ L) is added directly to each well and plates are incubated for 4 h in a humidified 5% CO ₂ atmosphere at 37°C. Absorbance is measured with a PowerWave XS2 Microplate Spectrophotometer at 490 nm and the results are expressed as percentage changes from the basal condition using four to five culture wells for each experimental treatment. In some experiments, HL60 cells are cultured with 100 nM ATRA and 1 μ M dasatinib for 4 days, and 10 μ M radotinib is added to each group according to the planned schedule(Only for Reference)
---------------	---

Solubility Information

Solubility	Ethanol: < 1 mg/mL (insoluble or slightly soluble), DMSO: 13.1 mg/mL (24.69 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.885 mL	9.4251 mL	18.8501 mL
5 mM	0.377 mL	1.885 mL	3.770 mL
10 mM	0.1885 mL	0.9425 mL	1.885 mL
50 mM	0.0377 mL	0.1885 mL	0.377 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

- Kim SH, et al. Haematologica. 2014, 99(7), 1191-1196.
Heo SK, et al. PLoS One. 2015, 10(6):e0129853.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481