


## NG-Nitroarginine methyl ester

## Chemical Properties

CAS No. : 50903-99-6  
 Formula: C<sub>7</sub>H<sub>15</sub>N<sub>5</sub>O<sub>4</sub>  
 Molecular Weight: 233.23  
 Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year  
 Actual storage temperature shall be subject to the COA.



## Biological Description

Description	NG-Nitroarginine methyl ester (ω-Nitroarginine methyl) is a non-selective nitric oxide synthase inhibitor. It has been utilized experimentally to induce hypertension.
Targets(IC50)	NO Synthase

## Solubility Information

Solubility	DMSO: 2.34 mg/mL (10.03 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	---

## Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	4.2876 mL	21.4381 mL	42.8761 mL
5 mM	0.8575 mL	4.2876 mL	8.5752 mL
10 mM	0.4288 mL	2.1438 mL	4.2876 mL
50 mM	0.0858 mL	0.4288 mL	0.8575 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

### Reference

Güney HZ, Görgün CZ, Tunçtan B, Uludağ O, Hodoğlugil U, Abacioğlu N, Zengil H. Circadian-rhythm-dependent effects of L-NG-nitroarginine methyl ester (L-NAME) on morphine-induced analgesia. *Chronobiol Int.* 1998 May;15(3):283-9.

Ke F, Kuang W, Hu X, et al. A novel vaccine targeting  $\beta$ 1-adrenergic receptor. *Hypertension Research.* 2023: 1-14.  
Tsuda K. Letter by Tsuda regarding article, "Plasminogen activator inhibitor-1 antagonist TM5441 attenuates Nw-nitro-L-arginine methyl ester-induced hypertension and vascular senescence". *Circulation.* 2014 Aug 26;130(9):e83.

Bulbul T, Akosman MS, Yilmaz O, Ulutas E, Bulbul A. Supplementary dietary nitric oxide donor (sodium nitroprusside) or inhibitor (NG-nitro-L-arginine methyl ester) depressed growth performance and ovarian primordial and primary follicles in Japanese quail (*Coturnix coturnix japonica*) in a dose-dependent manner. *Br Poult Sci.* 2015;56(1):113-20.

Shimizu M, Kochi T, Shirakami Y, Genovese S, Epifano F, Fiorito S, Mori T, Tanaka T, Moriwaki H. A newly synthesized compound, 4'-geranyloxyferulic acid-N(omega)-nitro-L-arginine methyl ester suppresses inflammation-associated colorectal carcinogenesis in male mice. *Int J Cancer.* 2014 Aug 15;135(4):774-84.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481