


Btk inhibitor 2

Chemical Properties

CAS No. :	1558036-85-3
Formula:	C ₂₄ H ₂₅ N ₅ O ₃
Molecular Weight:	431.49
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



Biological Description

Description	Btk Inhibitor 2 (BGB-3111 analog) is a BTK inhibitor.
Targets(IC50)	BTK

Solubility Information

Solubility	DMSO: 75 mg/mL (173.82 mM), Sonication and heating are recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 2 mg/mL (4.64 mM), Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.3176 mL	11.5878 mL	23.1755 mL
5 mM	0.4635 mL	2.3176 mL	4.6351 mL
10 mM	0.2318 mL	1.1588 mL	2.3176 mL
50 mM	0.0464 mL	0.2318 mL	0.4635 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Cecile M. Krejsa. Methods of Using BTK Inhibitors to Treat Dermatoses. US 20170224688 A1.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481