


Sulfasalazine

Chemical Properties

CAS No. :	599-79-1
Formula:	C ₁₈ H ₁₄ N ₄ O ₅ S
Molecular Weight:	398.39
Storage:	Keep away from direct sunlight, Keep away from moisture Powder: -20°C for 3 years In solvent: -80°C for 1 year <i>Actual storage temperature shall be subject to the COA.</i>



Biological Description

Description	Sulfasalazine (Azulfidine) is a synthetic salicylic acid derivative with affinity for elastin-containing connective tissues and formulated as a prodrug. Sulfasalazine induces iron death and inhibits NF- κ B, TGF- β and COX-2.
Targets(IC50)	Apoptosis, Ferroptosis, NF- κ B, Antibacterial, Antibiotic, Autophagy, COX
In vitro	<p>METHODS: Rat glioma cells F98 and human glioma cells U251 were treated with Sulfasalazine (200-400 μM) for 96 h. Cell viability was measured by MTT assay.</p> <p>RESULTS: Cell viability was significantly reduced in F98 at 200-400 μM Sulfasalazine concentration and U251 showed reduced cell viability at 400 μM Sulfasalazine. [1]</p> <p>METHODS: Mouse melanoma cells B16F10 and mouse embryonic fibroblasts MEF were treated with Sulfasalazine (10-1000 μM) for 24 h. Cellular ROS levels were measured by DCFDA staining.</p> <p>RESULTS: At lower Sulfasalazine concentrations (10-100 μM), no increase in intracellular ROS was observed. At higher concentrations of Sulfasalazine (800-1000 μM), there was an approximately 2.3-fold increase in intracellular ROS in B16F10 cells, while no increase in ROS was observed in MEF cells. [2]</p>
In vivo	<p>METHODS: To detect antitumor activity in vivo, Sulfasalazine (250 mg/kg) was administered intraperitoneally to C57BL/6N mice bearing B16F10 xenografts once daily for three days. Twenty-four hours after the third Sulfasalazine dose, local X-ray irradiation was applied to anesthetized tumor-bearing C57BL/6N mice at a dose of 4 Gy.</p> <p>RESULTS: Sulfasalazine alone did not significantly inhibit tumor growth; X-ray irradiation partially reduced tumor growth; the combination of Sulfasalazine and X-rays synergistically reduced tumor growth. [2]</p>
Cell Research	Sulfasalazine is dissolved in culture medium. SW620 cells are grown in Dulbecco's modified Eagle medium, supplemented with 10% heat-inactivated FCS, 2 mmol/liter glutamine, and 1% (wt/vol) penicillin/streptomycin. SW620 cells are transfected with the 3xlgkBLuc reporter construct. After 18 h, cells are incubated with either medium alone or with sulfasalazine (0.1, 0.2, 0.5, 1, 2, 5 mM) before stimulation with TNF α , LPS, or PMA. Luciferase assay is performed[1].

Solubility Information

A DRUG SCREENING EXPERT

Solubility	NaOH: 250 mg/mL (627.53 mM) DMSO: 200 mg/mL (502.02 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 20 mg/mL (50.2 mM), Suspension. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.5101 mL	12.5505 mL	25.101 mL
5 mM	0.502 mL	2.5101 mL	5.0202 mL
10 mM	0.251 mL	1.2551 mL	2.5101 mL
50 mM	0.0502 mL	0.251 mL	0.502 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

- Sehm T, et al. Sulfasalazine impacts on ferroptotic cell death and alleviates the tumor microenvironment and glioma-induced brain edema. *Oncotarget*. 2016 Jun 14;7(24):36021-36033.
- Ma R, Fang L, Chen L, et al. Ferroptotic stress promotes macrophages against intracellular bacteria. *Theranostics*. 2022, 12(5): 2266
- Nagane M, et al. Sulfasalazine, an inhibitor of the cystine-glutamate antiporter, reduces DNA damage repair and enhances radiosensitivity in murine B16F10 melanoma. *PLoS One*. 2018 Apr 12;13(4):e0195151.
- Lu Y, Yang Q, Su Y, et al. MYCN mediates TFRC-dependent ferroptosis and reveals vulnerabilities in neuroblastoma. *Cell Death & Disease*. 2021, 12(6): 1-14.
- Ni H, Ruan G, Sun C, et al. Tanshinone IIA inhibits gastric cancer cell stemness through inducing ferroptosis. *Environmental Toxicology*. 2021
- Chung WJ, et al. *J Neurochem*, 2009, 110(1), 182-193.
- Wang Y, Wang S, Xin Y, et al. Hydrogen sulfide alleviates the anxiety-like and depressive-like behaviors of type 1 diabetic mice via inhibiting inflammation and ferroptosis. *Life Sciences*. 2021: 119551.
- Wang Y, Wang S, Xin Y, et al. Hydrogen sulfide alleviates the anxiety-like and depressive-like behaviors of type 1 diabetic mice via inhibiting inflammation and ferroptosis[J]. *Life Sciences*. 2021: 119551.
- Lu Y, Yang Q, Su Y, et al. MYCN mediates TFRC-dependent ferroptosis and reveals vulnerabilities in neuroblastoma [J]. *Cell Death & Disease*. 2021, 12(6): 1-14.
- Cao Y, Wang Y, Li Y, et al. Angelic acid triggers ferroptosis in colorectal cancer cells via targeting and impairing NRF2 protein stability. *Journal of Natural Medicines*. 2024: 1-13.
- Hu C, Chen Y, Jin T, et al. A derivative of tanshinone IIA and salvianone, 15a, inhibits inflammation and alleviates DSS-induced colitis in mice by direct binding and inhibition of RIPK2. *Acta Pharmacologica Sinica*. 2024: 1-15.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481