

Sulfo-Cy3 NHS ester

Chemical Properties

CAS No. :	146368-16-3
Formula:	C35H41N3O10S2
Molecular Weight:	727.844
Storage:	Keep away from direct sunlight, Store at low temperature Store at -20°C <small>Actual storage temperature shall be subject to the COA.</small>

Biological Description

Description	Sulfo-Cy3 NHS ester is a sulphonated CY3 derivative and fluorophore dye with an Ex/Em of 550/570 nm. Sulfo-Cy3 NHS ester is an amine-reactive fluorescent probe commonly employed for labelling nucleic acids and proteins.
Targets(IC50)	Others
In vitro	<p>Reference for the usage of CY3-SE:</p> <p>1. Protein Preparation It is recommended to prepare the protein (antibody) solution at a concentration of 2 mg/mL, with a pH of 8.5 ± 0.5. If the pH is below 8.0, adjust it using 1 M sodium bicarbonate. Note: If the protein concentration is below 2 mg/mL, the labeling efficiency will significantly decrease. For optimal labeling efficiency, the recommended final protein concentration range is 2-10 mg/mL. The protein buffer must not contain primary amines (e.g., Tris or glycine) or ammonium ions, as these will affect labeling efficiency.</p> <p>2. Dye Preparation Dissolve CY3-SE dye in anhydrous DMSO to prepare a 10 mg/mL stock solution—Aliquot and store at -20°C or -80°C, protected from light.</p> <p>3. Dye Quantity Calculation The required amount of CY3-SE dye for the labeling reaction depends on the quantity of protein to be labeled. The optimal molar ratio of CY3-SE dye to protein is approximately 10. Example: If the protein to be labeled is 500 μL of 2 mg/mL IgG (MW = 150,000), dissolve 1 mg of CY3-SE dye in 100 μL DMSO. The required volume of CY3-SE is 4.88 μL, calculated as follows: 1) $\text{mmol (IgG)} = \text{mg/mL (IgG)} \times \text{mL (IgG)} / \text{MW (IgG)} = 2 \text{ mg/mL} \times 0.5 \text{ mL} / 150,000$ $\text{mg/mmol} = 6.7 \times 10^{-6} \text{ mmol}$ 2) $\text{mmol (CY3-SE)} = \text{mmol (IgG)} \times 10 = 6.7 \times 10^{-6} \text{ mmol} \times 10 = 6.7 \times 10^{-5} \text{ mmol}$ 3) $\mu\text{L (CY3-SE)} = \text{mmol (CY3-SE)} \times \text{MW (CY3-SE)} / \text{mg}/\mu\text{L (CY3-SE)} = 6.7 \times 10^{-5} \text{ mmol} \times 727.84 \text{ mg/mmol} / 0.01 \text{ mg}/\mu\text{L} = 4.88 \mu\text{L (CY3-SE)}$</p> <p>4. Labeling Reaction Slowly add the corresponding volume of 10 mg/mL CY3-SE dye to 0.5 mL of the protein sample solution. Mix gently, then briefly centrifuge to collect the sample at the bottom of the reaction tube. Incubate the reaction tube in the dark at room temperature with</p>

A DRUG SCREENING EXPERT

In vitro	gentle shaking for 60 minutes, inverting the tube several times every 10-15 minutes. 5. Protein Purification Purify the dye-protein conjugate using an appropriate method. The above information is based on published literature. Experimental procedures should be appropriately modified to meet specific research demands.
----------	---

Solubility Information

Solubility	DMSO: 40.00 mg/mL (54.96 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween-80+45% Saline: 2.00 mg/mL (2.75 mM),Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.3739 mL	6.8696 mL	13.7393 mL
5 mM	0.2748 mL	1.3739 mL	2.7479 mL
10 mM	0.1374 mL	0.687 mL	1.3739 mL
50 mM	0.0275 mL	0.1374 mL	0.2748 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Reid DA, et al. Organization and dynamics of the nonhomologous end-joining machinery during DNA double-strand break repair. Proc Natl Acad Sci U S A. 2015 May 19;112(20):E2575-84.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481