


Isoorientin

Chemical Properties

CAS No. :	4261-42-1
Formula:	C ₂₁ H ₂₀ O ₁₁
Molecular Weight:	448.38
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



Biological Description

Description	Isoorientin (Homoorientin) has antioxidant activity.
Targets(IC50)	Antioxidant, COX

Solubility Information

Solubility	DMSO: 125 mg/mL (278.78 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+90% Saline: < 10 mg/mL (22.3 mM), Lower concentrations may be soluble, but exact solubility limit is unknown. 10% DMSO+40% PEG300+5% Tween 80+45% Saline: 10 mg/mL (22.3 mM), Solution. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.2303 mL	11.1513 mL	22.3025 mL
5 mM	0.4461 mL	2.2303 mL	4.4605 mL
10 mM	0.223 mL	1.1151 mL	2.2303 mL
50 mM	0.0446 mL	0.223 mL	0.4461 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Cao J , Xia X , Dai X , et al. Flavonoids profiles, antioxidant, acetylcholinesterase inhibition activities of extract from *Dryothyrum boryanum* (Willd.) Ching[J]. *Food and Chemical Toxicology*, 2013, 55(Complete):121-128.

Jang S A, Hwang Y H, Kim T, et al. Anti-Osteoporotic and Anti-Adipogenic Effects of the Water Extract of *Drynaria roosii* Nakaike in Ovariectomized Mice Fed a High-Fat Diet. *Molecules*. 2019, 24(17): 3051

Characterization of a Pyranose Oxidase/C-Glycoside Oxidase from *Microbacterium* sp. 3H14, Belonging to the Unexplored Clade II of Actinobacterial POx/CGOx

Jang S A, Hwang Y H, Kim T, et al. Anti-Osteoporotic and Anti-Adipogenic Effects of the Water Extract of *Drynaria roosii* Nakaike in Ovariectomized Mice Fed a High-Fat Diet[J]. *Molecules*. 2019, 24(17): 3051.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481