


## 6-Thioguanine

## Chemical Properties

CAS No. :	154-42-7
Formula:	C <sub>5</sub> H <sub>5</sub> N <sub>5</sub> S
Molecular Weight:	167.19
Storage:	Powder: -20°C for 3 years   In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



## Biological Description

Description	6-Thioguanine (2-Amino-6-purinethiol) is an antineoplastic compound which also has antimetabolite action. The drug is used in the therapy of acute leukemia.
Targets(IC50)	Apoptosis,Endogenous Metabolite,Autophagy,DNA Methyltransferase,DUB,SARS-CoV
In vitro	6-Thioguanine selectively kills BRCA2-deficient tumors in xenograft models as effectively as PARP inhibitors. It eradicates cells and tumors that have developed resistance to PARP inhibitors or cisplatin through genetic reversal via the BRCA2 gene. Additionally, 6-Thioguanine is effective in eliminating PARP inhibitor-resistant tumors with BRCA1 deficiencies.
In vivo	6-Thioguanine is integrated into DNA, differing from typical DNA bases as it serves as a potent UVA chromophore with a peak absorbance at 342 nm. Its incorporation alters topoisomerase II-induced DNA cleavage both in the presence and absence of etoposide. Additionally, 6-Thioguanine induces similar apoptotic responses in mismatch repair-proficient and -deficient HCT116 and HeLa cells.

## Solubility Information

Solubility	DMSO: 62.5 mg/mL (373.83 mM),Sonication is recommended. H <sub>2</sub> O: < 1 mg/mL (insoluble or slightly soluble), Ethanol: < 1 mg/mL (insoluble or slightly soluble), (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 1 mg/mL (5.98 mM),Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

### Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	5.9812 mL	29.9061 mL	59.8122 mL
5 mM	1.1962 mL	5.9812 mL	11.9624 mL
10 mM	0.5981 mL	2.9906 mL	5.9812 mL
50 mM	0.1196 mL	0.5981 mL	1.1962 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

### Reference

Krynetskaia NF, et al. *FASEB J*, 2000, 14(14), 2339-2344.

Łagosz-Ćwik K, Melnykova M, Nieboga E, et al. Mapping of DNA methylationsensitive cellular processes in gingival and periodontal ligament fibroblasts in the context of periodontal tissue homeostasis. *Frontiers in Immunology*. 2023, 14.

Li M, Bei Z C, Yuan Y, et al. In-cell bioluminescence resonance energy transfer (BRET)-based assay uncovers ceritinib and CA-074 as SARS-CoV-2 papain-like protease inhibitors. *Journal of Enzyme Inhibition and Medicinal Chemistry*. 2024, 39(1): 2387417.

Marathias VM, et al. *Nucleic Acids Res*, 1999, 27(14), 2860-2867.

Yamane K, et al. *Clin Cancer Res*, 2005, 11(6), 2355-2363.

Brem R, et al. *Photochem Photobiol*, 2012, 88(1), 5-13.

Issaeva N, et al. *Cancer Res*, 2010, 70(15), 6268-6276.

LaDuke KE, et al. Effects of azathioprine, 6-mercaptopurine, and 6-thioguanine on canine primary hepatocytes. *Am J Vet Res*. 2015 Jul;76(7):649-55.

Chuang SJ, et al. 6-Thioguanine is a noncompetitive and slow binding inhibitor of human deubiquitinating protease USP2. *Sci Rep*. 2018 Feb 15;8(1):3102.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E\_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481