


## Methapyrilene hydrochloride

## Chemical Properties

CAS No. :	135-23-9
Formula:	C <sub>14</sub> H <sub>20</sub> ClN <sub>3</sub> S
Molecular Weight:	297.85
Storage:	Powder: -20°C for 3 years   In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



## Biological Description

Description	Methapyrilene hydrochloride is a biochemical.
Targets(IC50)	Others,Histamine Receptor

## Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.3574 mL	16.787 mL	33.5739 mL
5 mM	0.6715 mL	3.3574 mL	6.7148 mL
10 mM	0.3357 mL	1.6787 mL	3.3574 mL
50 mM	0.0671 mL	0.3357 mL	0.6715 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

### Reference

- Kindrat I, Dreval K, Shpileva S, Tryndyak V, de Conti A, Mudalige TK, Chen T, Erstenyuk AM, Beland FA, Pogribny IP. Effect of methapyrilene hydrochloride on hepatic intracellular iron metabolism in vivo and in vitro. *Toxicol Lett.* 2017 Nov 5;281:65-73. doi: 10.1016/j.toxlet.2017.09.011. Epub 2017 Sep 19. PubMed PMID: 28935588.
- Inoue K, Ochi A, Koda A, Wako Y, Kawasako K, Doi T. The 14-day repeated dose liver micronucleus test with methapyrilene hydrochloride using young adult rats. *Mutat Res Genet Toxicol Environ Mutagen.* 2015 Mar;780-781:123-7. doi: 10.1016/j.mrgentox.2014.04.004. Epub 2014 Apr 24. PubMed PMID: 24768639.
- Doktorova TY, Ates G, Vinken M, Vanhaecke T, Rogiers V. Way forward in case of a false positive in vitro genotoxicity result for a cosmetic substance? *Toxicol In Vitro.* 2014 Feb;28(1):54-9. doi: 10.1016/j.tiv.2013.09.022. Epub 2013 Oct 2. PubMed PMID: 24095862.
- Yafune A, Taniai E, Morita R, Akane H, Kimura M, Mitsumori K, Shibutani M. Immunohistochemical cellular distribution of proteins related to M phase regulation in early proliferative lesions induced by tumor promotion in rat two-stage carcinogenesis models. *Exp Toxicol Pathol.* 2014 Jan;66(1):1-11. doi: 10.1016/j.etp.2013.07.001. Epub 2013 Jul 23. PubMed PMID: 23890812.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481