

Cirsimaritin

Chemical Properties


CAS No. : 6601-62-3

Formula: C₁₇H₁₄O₆

Molecular Weight: 314.29

Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year

Actual storage temperature shall be subject to the COA.



Biological Description

Description	Cirsimaritin exhibits antibacterial, anti-inflammation, anti-tumor, antioxidant properties and renal protection. Cirsimaritin binds weakly to the benzodiazepine site on GABAA receptors.
Targets(IC50)	GABA Receptor
In vitro	Cirsimaritin inhibits the growth of tumor cells and induced mitochondrial apoptosis in human gallbladder carcinoma cell line (GBC-SD), it triggers endoplasmic reticulum (ER) stress and down-regulates the phosphorylation of Akt [1]. Cirsimaritin increases tyrosinase activity and melanin content in murine B16F10 melanoma cells by activation of CREB as well as upregulation of MITF and tyrosinase expression in a dose-dependent manner [2].
In vivo	Cirsimaritin significantly reduces anxiety in mice. The high-affinity benzodiazepine binding site is not involved in the anxiolytic activity induced by these compounds. Cirsimaritin does not decrease by co-administration of flumazenil (2.5 mg/kg) with salvigenin, rosmanol and cirsimaritin (10 mg/kg)[3].

Solubility Information

Solubility	DMSO: Soluble, (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	---

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.1818 mL	15.9089 mL	31.8177 mL
5 mM	0.6364 mL	3.1818 mL	6.3635 mL
10 mM	0.3182 mL	1.5909 mL	3.1818 mL
50 mM	0.0636 mL	0.3182 mL	0.6364 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Quan Z, et al. Reactive oxygen species-mediated endoplasmic reticulum stress and mitochondrial dysfunction contribute to cirsimaritin-induced apoptosis in human gallbladder carcinoma GBC-SD cells. *Cancer Lett.* 2010 Sep 28;295(2):252-9.

Kim HJ, et al. Melanogenesis-inducing effect of cirsimaritin through increases in microphthalmia-associated transcription factor and tyrosinase expression. *Int J Mol Sci.* 2015 Apr 20;16(4):8772-88.

Abdelhalim A, et al. Antidepressant, Anxiolytic and Antinociceptive Activities of Constituents from *Rosmarinus Officinalis*. *J Pharm Pharm Sci.* 2015;18(4):448-59.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481