

PF-4800567

## Chemical Properties

CAS No. : 1188296-52-7


Formula: C<sub>17</sub>H<sub>18</sub>ClN<sub>5</sub>O<sub>2</sub>

Molecular Weight: 359.81

Storage: Keep away from direct sunlight, Store under nitrogen,  
Store at low temperature

Powder: -20°C for 3 years | In solvent: -80°C for 1 year

Actual storage temperature shall be subject to the COA.



## Biological Description

Description	PF-4800567 is a selective inhibitor of casein kinase 1ε (CK1ε; IC <sub>50</sub> = 32 nM) with greater than 20-fold selectivity over CK1δ.
Targets(IC <sub>50</sub> )	Casein Kinase
In vitro	PF-4800567 shows inhibitory activity against CK1ε and CK1δ in whole cells, with IC <sub>50</sub> s of 2.65 and 20.38 μM, respectively. PF-4800567 (0.01-10 μM) blocks CK1ε-mediated PER3 nuclear localization mediated by CK1ε and suppresses PER2 degradation at 1 μM. In addition, PF-4800567 (32 nM) has little effect on the circadian clock [1].
In vivo	PF-4800567 (100 mg/kg, s.c.) is quickly absorbed and distributed in the plasma and brain of mice [1].

## Solubility Information

Solubility	DMSO: 62.5 mg/mL (173.7 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+90% Saline: < 6.25 mg/mL (17.37 mM), Lower concentrations may be soluble, but exact solubility limit is unknown. 10% DMSO+40% PEG300+5% Tween 80+45% Saline: 6.25 mg/mL (17.37 mM), Solution. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

### Preparing Stock Solutions

---

	<b>1mg</b>	<b>5mg</b>	<b>10mg</b>
1 mM	2.7792 mL	13.8962 mL	27.7924 mL
5 mM	0.5558 mL	2.7792 mL	5.5585 mL
10 mM	0.2779 mL	1.3896 mL	2.7792 mL
50 mM	0.0556 mL	0.2779 mL	0.5558 mL

---

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

### Reference

Walton KM, et al. Selective inhibition of casein kinase 1 epsilon minimally alters circadian clock period. J Pharmacol Exp Ther. 2009 Aug;330(2):430-9.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481