


4-Hydroxyantipyrine

Chemical Properties

CAS No. :	1672-63-5
Formula:	C ₁₁ H ₁₂ N ₂ O ₂
Molecular Weight:	204.23
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



Biological Description

Description	4-Hydroxyantipyrine is formed during oxidative deamination of aminopyrine.
Targets(IC50)	COX, Drug Metabolite

Solubility Information

Solubility	DMSO: 45 mg/mL (220.34 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 2 mg/mL (9.79 mM), Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	4.8964 mL	24.4822 mL	48.9644 mL
5 mM	0.9793 mL	4.8964 mL	9.7929 mL
10 mM	0.4896 mL	2.4482 mL	4.8964 mL
50 mM	0.0979 mL	0.4896 mL	0.9793 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Brune K, Neubert A. Clin Exp Rheumatol. 2001 Nov-Dec;19(6 Suppl 25):S51-7.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481