


Exendin-4

Chemical Properties

CAS No. :	141758-74-9
Formula:	C184H282N50O60S
Molecular Weight:	4186.57
Storage:	Keep away from moisture, Store at low temperature Powder: -20°C for 3 years In solvent: -80°C for 1 year <i>Actual storage temperature shall be subject to the COA.</i>



Biological Description

Description	Exendin-4 (Exenatide) is a glucagon-like peptide-1 receptor (GLP-1) agonist (IC50: 3.22 nM). Exenatide is a 39 amino acid peptide. Compared to GLP-1, exenatide has a longer half-life of 2.4 hours.
Targets(IC50)	Glucagon Receptor
In vitro	In HUVECs, exendin-4 dose-dependently significantly increases NO production, eNOS phosphorylation and GTPCH1 level[2]. Exendin-4 shows cytotoxic effects to MCF-7 breast cancer cells (IC50 5 µM) at 48 hours [3].
In vivo	In ob/ob mice, the treatment of exendin-4 improve serum ALT and reduce serum glucose, insulin levels and calculated HOMA scores compared with control. In the final 4 weeks of the study period, exendin-4-treated ob/ob mice sustain an obvious reduction in the net weight gain[4]. Animals treated with exendin-4 have more pyknotic nuclei, more pancreatic acinar inflammation and weigh significantly less than control rats[5]. Exenatide leads to dose-dependent relaxation of rat thoracic aorta, which is evoked via the GLP-1 receptor and is mediated mainly by H2S but also by CO and NO[6].

Solubility Information

Solubility	DMSO: 10 mM, Sonication is recommended. H2O: 1.23 mg/mL (0.29 mM), Sonication and heating are recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 2 mg/mL (0.48 mM), Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	0.2389 mL	1.1943 mL	2.3886 mL
5 mM	0.0478 mL	0.2389 mL	0.4777 mL
10 mM	0.0239 mL	0.1194 mL	0.2389 mL
50 mM	0.0048 mL	0.0239 mL	0.0478 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

- Doyle ME, et al. The importance of the nine-amino acid C-terminal sequence of exendin-4 for binding to the GLP-1 receptor and for biological activity. *Regul Pept.* 2003 Jul 15;114(2-3):153-8.
- Huang L, Ma R, Lin T, et al. Glucagon-like peptide-1 receptor pathway inhibits extracellular matrix production by mesangial cells through store-operated Ca²⁺ channel. *Experimental Biology and Medicine.* 2019: 1535370219876531.
- Ren W, Chen J, Wang W, et al. Sympathetic nerve-enteroendocrine L cell communication modulates GLP-1 release, brain glucose utilization, and cognitive function. *Neuron.* 2024
- Wei R, et al. Exenatide exerts direct protective effects on endothelial cells through the AMPK/Akt/eNOS pathway in a GLP-1 receptor-dependent manner. *Am J Physiol Endocrinol Metab.* 2016 Jun 1; 310(11):E947-57.
- Ding X, et al. Exendin-4, a glucagon-like protein-1 (GLP-1) receptor agonist, reverses hepatic steatosis in ob/obmice. *Hepatology.* 2006 Jan;43(1):173-81.
- Nachnani JS, et al. Biochemical and histological effects of exendin-4 (exenatide) on the rat pancreas. *Diabetologia.* 2010 Jan;53(1):153-9.
- Sélley, E., Kun, S., Szijártó, I., Laczy, B., Kovács, T., & Fülöp, F. et al. (2014). Exenatide induces aortic vasodilation increasing hydrogen sulphide, carbon monoxide and nitric oxide production. *Cardiovascular Diabetology*, 13(1), 69. doi: 10.1186/1475-2840-13-69
- Fidan-Yaylall G, et al. Antidiabetic exendin-4 activates apoptotic pathway and inhibits growth of breast cancer cells. *Tumour Biol.* 2016 Feb;37(2):2647-53.
- Huang L, Ma R, Lin T, et al. Glucagon-like peptide-1 receptor pathway inhibits extracellular matrix production by mesangial cells through store-operated Ca²⁺ channel[J]. *Experimental Biology and Medicine.* 2019: 1535370219876531.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481