

Venetoclax

Chemical Properties

CAS No. : 1257044-40-8

Formula: C₄₅H₅₀ClN₇O₇S


Molecular Weight: 868.44

Storage:

Keep away from direct sunlight, Keep away from moisture, Store at low temperature

Powder: -20°C for 3 years | In solvent: -80°C for 1 year

Actual storage temperature shall be subject to the COA.



Biological Description

Description	Venetoclax (ABT-199) is a Bcl-2 inhibitor (K _i <0.01 nM) with potent, selective, and orally active properties. Venetoclax has a 3-order-of-magnitude lower affinity for Bcl-xL and Bcl-W (K _i =48/245 nM). Venetoclax induces autophagy and apoptosis.
Targets(IC50)	Bcl-2 Family, Autophagy
In vitro	<p>METHODS: Eleven human T-cell acute lymphoblastic leukemia cells T-ALL were treated with Venetoclax (0-12 μM) for 48 h. Cell viability was measured by Celltiter-Glo Luminescent Cell Viability Assay.</p> <p>RESULTS: The IC₅₀ values of Venetoclax against 11 T-ALL species ranged from 0.2-10 μM. [1]</p> <p>METHODS: Human acute lymphoid leukemia cells RS4;11 were incubated with Venetoclax (0.01-5 μM) for 3.5 h. Caspase-3/7 activity was assessed using the Caspase-GLO kit.</p> <p>RESULTS: Venetoclax induced the activation of caspases, which is one of the characteristics of apoptosis. [2]</p> <p>METHODS: Human primary HCL leukemia cells were treated with Venetoclax (0.1-1 μM) for 24 h. Cell death was detected using Flow Cytometry method.</p> <p>RESULTS: Venetoclax significantly increased cell death in HCL cells in a dose-dependent manner. [3]</p>
In vivo	<p>METHODS: To test the antitumor activity in vivo, Venetoclax (100 mg/kg in 60% PG+30% PEG 400+10% ethanol) was orally administered once daily for twenty-one days to C.B-17 SCID-beige mice bearing human diffuse large B-cell lymphoma Toledo.</p> <p>RESULTS: Venetoclax significantly inhibited the growth of Toledo tumor (TGI_{max}=93%, TGD=220%). [2]</p> <p>METHODS: To assay anti-tumor activity in vivo, Venetoclax (50 mg/kg in 10% ethanol+30% PEG 400+60% Phosal 50PG, administered orally once daily) and anti-PD-1 (10 mg/kg in PBS, administered intraperitoneally three times every four days) were administered to C57BL/6 mice harboring mouse colorectal carcinoma tumor MC38 for fourteen days.</p> <p>RESULTS: Venetoclax enhances the antitumor efficacy of immune checkpoint inhibitors (ICIs) and increases PD-1+ T effector memory cells. Venetoclax does not impair human T-cell function in response to antigenic stimulation in vitro, nor does it antagonize anti-</p>

In vivo	PD-1-induced T-cell activation. [4]
Kinase Assay	The equilibrium binding experiments of fluorescent peptides to Bcl-xL protein were performed in an Analyst 96-well plate reader under the following conditions: each individual well in a 96-well assay plate contained 5 μ l DMSO, 15 nM fluorescent peptide, and increasing concentrations (from 0 to 2.24 μ M) of Bcl-xL protein in assay buffer in a final volume of 125 μ l. The plate was mixed on a shaker for 1 min and incubated at room temperature for an additional 15 min. The polarization in millipolarization units (mP) was measured at room temperature with an excitation wavelength at 485 nm and an emission wavelength at 530 nm. For assay stability testing, a plate containing a binding experiment was measured at different times over a 24-h period. Between each reading, the plate was covered with parafilm to prevent any solution evaporation. To determine the effect of DMSO on the assay, binding experiments were performed under conditions similar to those described above except that the amount of DMSO was varied from 0 to 4 to 8%. All experimental data were analyzed using Prism 3.0 software and Kd values were generated by fitting the experimental data using a sigmoidal dose-response nonlinear regression model [1].
Cell Research	RS4;11 cells were seeded at 50,000 per well in 96-well plates and treated with compounds diluted in half-log steps starting at 1 μ M and ending at 0.00005 μ M. All other leukemia and lymphoma cell lines were seeded at 15,000–20,000 cells per well in the appropriate medium and incubated with ABT-199 or navitoclax for 48 h. Effects on proliferation were determined using Cell TiterGlo reagent. EC50 values were determined by nonlinear regression analysis of the concentration-response data. Mouse FL5.12-BCL-2 and FL5.12-BCL-XL cells were propagated and assessed as described previously. Bak ^{+/+} Bax ^{+/+} double knockout mouse embryonic fibroblasts were seeded into 96-well microtiter plates at 5,000 cells per well in DMEM supplemented with 10% FBS. ABT-199 in the same culture medium was added in half-log dilutions starting at 5 μ M. The cells were then incubated at 37 °C (5% CO ₂) for 48 h, and the effects on proliferation were determined using Cell TiterGlo reagent according to the manufacturer's instructions [1].
Animal Research	Female C.B-17 SCID mice (DoHH2 and Granta-519 xenografts) and female C.B-17 SCID-beige mice (RS4;11 and Toledo xenografts) were inoculated with 1 \times 10 ⁶ (DoHH2) or 5 \times 10 ⁶ (Granta-519, Toledo and RS4;11) cells subcutaneously in the right flank. The inoculation volume (0.2 ml) comprised a 50:50 mixture of cells in growth media and Matrigel. Electronic calipers were used to measure the length and width of each tumor 2–3 times per week. Tumor volume was estimated by applying the following equation: volume = length \times width ² /2. When tumors reached approximately 220 mm ³ , mice were size matched (day 0) into treatment and control groups. All xenograft trials were conducted using ten mice per group, and all mice were ear tagged and monitored individually throughout the studies. ABT-199 was formulated for oral dosing in 60% phosal 50 propylene glycol (PG), 30% polyethylene glycol (PEG) 400 and 10% ethanol, and bendamustine and rituximab were formulated in accordance with the manufacturer's instructions. ABT-199 was delivered approximately 2 h before bendamustine or bendamustine plus rituximab. TGImax was calculated as the greatest treatment response using the following equation: TGImax = (1 - mean tumor volume of the treated group/mean tumor volume of the vehicle control group) \times 100. The TGD (%) was determined as the percentage increase of the median time period for the treatment group to reach an arbitrary tumor volume of 1,000 mm ³ relative to the vehicle control group. A complete tumor regression response was the portion of the population with tumors \leq 25 mm ³ for at least three consecutive measurements [1].

Solubility Information

Solubility	H2O: < 1 mg/mL (insoluble or slightly soluble), Ethanol: < 1 mg/mL (insoluble or slightly soluble), DMSO: 257.5 mg/mL (296.51 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 10 mg/mL (11.51 mM),Suspension. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.1515 mL	5.7575 mL	11.5149 mL
5 mM	0.2303 mL	1.1515 mL	2.303 mL
10 mM	0.1151 mL	0.5757 mL	1.1515 mL
50 mM	0.023 mL	0.1151 mL	0.2303 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

- Peirs S, et al. ABT-199 mediated inhibition of BCL-2 as a novel therapeutic strategy in T-cell acute lymphoblastic leukemia. *Blood*. 2014 Dec 11;124(25):3738-47.
- Lima K, Pereira-Martins D A, de Miranda L B L, et al. The PIP4K2 inhibitor THZ-P1-2 exhibits antileukemia activity by disruption of mitochondrial homeostasis and autophagy. *Blood cancer journal*. 2022, 12(11): 1-11.
- Lee J B, Khan D H, Hurren R, et al. Venetoclax enhances T cell-mediated anti-leukemic activity by increasing ROS production. *Blood*. 2021 Jul 22;138(3):234-245. doi: 10.1182/blood.2020009081.
- Capoci I R G, Faria D R, Sakita K M, et al. Repurposing approach identifies new treatment options for invasive fungal disease. *Bioorganic Chemistry*. 2019 Mar;84:87-97
- Hirakata, Camila, et al. Targeting glioma cells by antineoplastic activity of reversine. *Oncology Letters*. 22.2 (2021): 1-9
- e Silva, Catarina Sofia Mateus Reis, et al. Embelin potentiates venetoclax-induced apoptosis in acute myeloid leukemia cells. *Toxicology in Vitro*. (2021): 105207.
- Lima K, Carvalho M F L, Pereira-Martins D A, et al. Pharmacological Inhibition of PIP4K2 Potentiates Venetoclax-Induced Apoptosis in Acute Myeloid Leukemia. *International Journal of Molecular Sciences*. 2023, 24(23): 16899.
- Diwan R, Bhatt H N, Dong R, et al. Cell selective BCL-2 inhibition enabled by lipid nanoparticles alleviates lung fibrosis. *Journal of Controlled Release*. 2024, 370: 421-437.
- Small S H, Perez R E, Beauchamp E M, et al. Targeting SLFN11-Regulated Pathways Restores Chemotherapy Sensitivity in AML. *Blood Neoplasia*. 2024: 100037.
- Zhao L, Sun X, Hou C, et al. CPNE7 promotes colorectal tumorigenesis by interacting with NONO to initiate ZFP42 transcription. *Cell Death & Disease*. 2024, 15(12): 896.
- Luo X, Linghu M, Zhou X, et al. Merestinib inhibits cuproptosis by targeting NRF2 to alleviate acute liver injury. *Free Radical Biology and Medicine*. 2025
- Souers AJ, et al. ABT-199, a potent and selective BCL-2 inhibitor, achieves antitumor activity while sparing platelets. *Nat Med*. 2013 Feb;19(2):202-8.
- Pan G, Zhong M, Yao J, et al. Orelabrutinib and venetoclax synergistically induce cell death in double-hit lymphoma by interfering with the crosstalk between the PI3K/AKT and p38/MAPK signaling. *Journal of Cancer Research and Clinical Oncology*. 2022: 1-17.
- Gao P, Zhang W, Fang X, et al. Simultaneous quantification of venetoclax and voriconazole in human plasma by UHPLC-MS/MS and its application in acute myeloid leukemia patients. *Journal of Pharmaceutical and Biomedical Analysis*. 2023: 115279.
- Vereertbrugghen A, et al. In Vitro Sensitivity to Venetoclax and Microenvironment Protection in Hairy Cell Leukemia. *Front Oncol*. 2021 Jul 26;11:598319.
- Kohlhapp FJ, et al. Venetoclax Increases Intratumoral Effector T Cells and Antitumor Efficacy in Combination with Immune Checkpoint Blockade. *Cancer Discov*. 2021 Jan;11(1):68-79.
- Li Z, Pan G, Zhong M, et al. High-Throughput Drug Screen for Potential Combinations With Venetoclax Guides the Treatment of Transformed Follicular Lymphoma. *International Journal of Toxicology*. 2023: 10915818231178693.
- Capoci I R G, Faria D R, Sakita K M, et al. Repurposing approach identifies new treatment options for invasive fungal disease[J]. *Bioorganic chemistry*. 2019 Mar;84:87-97.
- e Silva, Catarina Sofia Mateus Reis, et al. Embelin potentiates venetoclax-induced apoptosis in acute myeloid leukemia cells. *Toxicology in Vitro*. (2021): 105207.
- Takei H, Coelho-Silva J L, Leal C T, et al. Suppression of multiple anti-apoptotic BCL2 family proteins recapitulates the effects of JAK2 inhibitors in JAK2V617F driven myeloproliferative neoplasms. *Cancer Science*. 2022, 113(2): 597.
- Suarez M, Blyth G T, Mina A A, et al. Inhibitory effects of Tomivosertib in acute myeloid leukemia. *Oncotarget*. 2021, 12(10): 955.
- Tang J, Yao C, Liu Y, et al. Arsenic trioxide induces expression of BCL-2 expression via NF- κ B and p38 MAPK signaling pathways in BEAS-2B cells during apoptosis. *Ecotoxicology and Environmental Safety*. 2021, 222: 112531.
- Tan M, Ren F, Yang X. Anti-HBV therapeutic potential of small molecule 3, 5, 6, 7, 3', 4'-Hexamethoxyflavone in vitro and in vivo. *Virology*. 2021

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481