


## 2'-Deoxyadenosine

## Chemical Properties

|                   |   |
|-------------------|---|
| CAS No. :         | 958-09-8  |
| Formula:          | C10H13N5O3  |
| Molecular Weight: | 251.24  |
| Storage:          | Powder: -20°C for 3 years   In solvent: -80°C for 1 year<br>Actual storage temperature shall be subject to the COA. |



## Biological Description

|               |  |
|---------------|--|
| Description   | 2'-Deoxyadenosine (NSC-141848) is the DNA nucleoside A. It pairs with deoxythymidine (T) in double-stranded DNA. |
| Targets(IC50) | Apoptosis, Nucleoside Antimetabolite/Analog, Caspase, Endogenous Metabolite                                      |

## Solubility Information

|                     |   |
|---------------------|---|
| Solubility          | DMSO: 2.52 mg/mL (10.03 mM), Sonication is recommended.<br>(< 1 mg/ml refers to the product slightly soluble or insoluble)  |
| In vivo Formulation | 10% DMSO+40% PEG300+5% Tween 80+45% Saline: 1 mg/mL (3.98 mM), Sonication is recommended.<br><i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i> |

## Preparing Stock Solutions

|       | 1mg       | 5mg        | 10mg       |
|-------|-----------|------------|------------|
| 1 mM  | 3.9803 mL | 19.9013 mL | 39.8026 mL |
| 5 mM  | 0.7961 mL | 3.9803 mL  | 7.9605 mL  |
| 10 mM | 0.398 mL  | 1.9901 mL  | 3.9803 mL  |
| 50 mM | 0.0796 mL | 0.398 mL   | 0.7961 mL  |

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

### Reference

- Murakami H, Aoyanagi T, Miki Y, Tomita H, Esaka Y, Inoue Y, Teshima N. Effects of hydrophilic monomers on sorptive properties of divinylbenzene-based reversed phase sorbents. *Talanta*. 2018 Aug 1;185:427-432. doi: 10.1016/j.talanta.2018.03.093. Epub 2018 Mar 30. PubMed PMID: 29759223.
- Markowitz M, Sarafianos SG. 4'-Ethynyl-2'-fluoro-2'-deoxyadenosine, MK-8591: a novel HIV-1 reverse transcriptase translocation inhibitor. *Curr Opin HIV AIDS*. 2018 Apr 24. doi: 10.1097/COH.0000000000000467. [Epub ahead of print] PubMed PMID: 29697468.
- Stegmüller S, Schrenk D, Cartus AT. Formation and fate of DNA adducts of alpha- and beta-asarone in rat hepatocytes. *Food Chem Toxicol*. 2018 Jun;116(Pt B):138-146. doi: 10.1016/j.fct.2018.04.025. Epub 2018 Apr 11. PubMed PMID: 29654846.
- Carlson ES, Upadhyaya P, Villalta PW, Ma B, Hecht SS. Analysis and Identification of 2'-Deoxyadenosine-Derived Adducts in Lung and Liver DNA of F-344 Rats Treated with the Tobacco-Specific Carcinogen 4-(Methylnitrosamino)-1-(3-pyridyl)-1-butanone and Enantiomers of its Metabolite 4-(Methylnitrosamino)-1-(3-pyridyl)-1-butanol. *Chem Res Toxicol*. 2018 Apr 19. doi: 10.1021/acs.chemrestox.8b00056. [Epub ahead of print] PubMed PMID: 29651838.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481