


Pacritinib

Chemical Properties

CAS No. :	937272-79-2
Formula:	C ₂₈ H ₃₂ N ₄ O ₃
Molecular Weight:	472.58
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



Biological Description

Description	Pacritinib (SB1518) (SB1518) is an effective and specific inhibitor of JAK2 and FLT3 (IC ₅₀ : 23/22 nM, in cell-free assays).
Targets(IC ₅₀)	FLT,JAK,Tyrosine Kinases
In vitro	Pacritinib is an effective inhibitor of both wild-type JAK2 and JAK2V617F mutant (IC ₅₀ : 19 nM). The IC ₅₀ s of Pacritinib are 50 nM for TYK2, 520 nM for JAK3 and 1280 nM for JAK1. Pacritinib effectively permeates cells to modulate signaling pathways downstream of JAK2, whether agonist activated or mutationally activated. Pacritinib induces apoptosis, cell cycle arrest and antiproliferative effects in JAK2WT- and JAK2V617F-dependent cells. Pacritinib inhibits cell proliferation of Karpas 1106P (IC ₅₀ : 348 nM) and Ba/F3-JAK2V617F (IC ₅₀ : 160 nM), respectively. Pacritinib inhibits endogenous colony growth derived from erythroid (IC ₅₀ : 63 nM) and myeloid progenitors(IC ₅₀ : 53 nM), respectively. [1] SB1518 also inhibits FLT3 and its mutant FLT3-D835Y (IC ₅₀ : 6 nM). Pacritinib inhibits FLT3 phosphorylation and downstream STAT, MAPK and PI3K signaling in FLT3-internal-tandem duplication (ITD), FLT3-wt cells and primary AML blast cells. Pacritinib treatment leads to a dose-dependent decrease of pFLT3, pSTAT5, pERK1/2 and pAkt in FLT3-ITD harboring MV4-11 cells with IC ₅₀ of 80, 40, 33 and 29 nM , respectively. Treatment of the primary AML blast cells with Pacritinib for 3 h leads to a dose-dependent decrease of pFLT3, pSTAT3 and pSTAT5 with an IC ₅₀ below 0.5 μM. Pacritinib induces apoptosis, cell cycle arrest and anti-proliferative effects in FLT3-mutant and FLT3-wt cells. Pacritinib inhibits cell proliferation of FLT3-ITD-harboring cells MV4-11 (IC ₅₀ : 47 nM) and primary AML blast (IC ₅₀ : 0.19-1.3 nM) cells.
In vivo	In JAK2V617F-dependent xenograft model, Pacritinib (150 mg/kg, p.o., q.d.) markedly ameliorates splenomegaly and hepatomegaly symptoms, with 60% normalization of spleen weight and 92% normalization of liver weight and is well tolerated without significant weight loss or any hematological toxicities, including thrombocytopenia and anemia. In JAK2V617F-dependent SET-2 xenograft model, Pacritinib dose-dependent inhibits tumor growth (40% for 75 mg/kg and 61% for 150 mg/kg).[1] Pacritinib treated once daily for 21 consecutive days, induces dose-dependent inhibition of tumor growth (38% for 25 mg/kg, 92% for 50 mg/kg and 121% for 100 mg/kg). Complete regression is observed in 3/10 and 8/8 mice for the 50 and 100 mg/kg/day groups, respectively.
Kinase Assay	kinase activity assays: All assays are carried out in 384-well white microtiter plates. Compounds are 4-fold serially diluted in 8 steps, starting from 10 μM. The reaction

Kinase Assay	mixture consisted of 25 μ L assay buffer (50 mM HEPES pH 7.5, 10 mM MgCl ₂ , 5 mM MnCl ₂ , 1 mM DTT, 0.1 mM Na ₃ VO ₄ , 5 mM β -glycerol phosphate). For FLT3 assays, the reaction contains 2.0 μ g/mL FLT3 enzyme, 5 μ M of poly(Glu,Tyr) substrate and 4 μ M of ATP. For JAK1 assays, the reaction contains 2.5 μ g/mL of JAK1 enzyme, 10 μ M of poly (Glu,Ala,Tyr) substrate and 1.0 μ M of ATP. For JAK2 assays, the reaction contained 0.35 μ g/mL of JAK2 enzyme, 10 μ M of poly (Glu,Ala,Tyr) substrate and 0.15 μ M of ATP. For JAK3 assays, the reaction contained 3.5 μ g/mL of JAK3 enzyme, 10 μ M of poly (Glu,Ala, Tyr) substrate and 6.0 μ M of ATP. For TYK2 assays, the reaction contained 2.5 μ g/mL of TYK2 enzyme, 10 μ M of poly (Glu,Ala,Tyr) substrate and 0.15 μ M of ATP. The reaction is incubated at room temperature for 2 h prior to addition of 13 μ L PKLight [®] detection reagent. After 10 min incubation luminescent signals are read on a multi-label plate reader.
Cell Research	Cells are seeded at 30-50% confluency in 96-well plates and are treated with different concentrations of compounds (in triplicate) for 48 h. Cell viability is monitored using the CellTiter-Glo assay.(Only for Reference)

Solubility Information

Solubility	H ₂ O: < 1 mg/mL (insoluble or slightly soluble), Ethanol: < 1 mg/mL (insoluble or slightly soluble), DMSO: 1.94 mg/mL (4.11 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 0.36 mg/mL (0.76 mM),Solution. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.116 mL	10.5802 mL	21.1604 mL
5 mM	0.4232 mL	2.116 mL	4.2321 mL
10 mM	0.2116 mL	1.058 mL	2.116 mL
50 mM	0.0423 mL	0.2116 mL	0.4232 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Hart S, et al. Leukemia, 2011, 25(11), 1751-1759.

Chen K Y, Krischuns T, Varga L O, et al. A highly sensitive cell-based luciferase assay for high-throughput automated screening of SARS-CoV-2 nsp5/3CLpro inhibitors. Antiviral Research. 2021

Hart S, et al. Blood Cancer J, 2011, 1(11), e44.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481