


Lurasidone hydrochloride

Chemical Properties

CAS No. : 367514-88-3
 Formula: C₂₈H₃₆N₄O₂S·HCl
 Molecular Weight: 529.14
 Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year
Actual storage temperature shall be subject to the COA.



Biological Description

Description	Lurasidone hydrochloride (Lurasidone HCl) is a thiazole derivative and atypical antipsychotic agent that functions as a dopamine D2 receptor antagonist; serotonin 5-HT ₂ receptor antagonist, serotonin 5-HT ₇ receptor antagonist, an antagonist of the adrenergic α _{2A} and α _{2C} receptors, as well as a partial serotonin 5-HT _{1A} receptor agonist. It is used in the treatment of schizophrenia and bipolar disorder.
Targets(IC50)	5-HT Receptor, Norepinephrine, Dopamine Receptor
In vitro	Lurasidone antagonizes dopamine-stimulated [³⁵ S]GTP γ S binding at human dopamine D _{2L} receptor in a concentration-dependent manner with a K _B value of 2.8 nM. Lurasidone antagonizes 5-HT-stimulated cAMP accumulation in the CHO/h5-HT ₇ cells with a K _B value of 2.6 nM. Lurasidone partially stimulates [³⁵ S]GTP γ S binding to the membrane preparation for human 5-HT _{1A} receptors with a maximum effect of 33%. Lurasidone dose-dependently increases the ratio of DOPAC/dopamine in rat frontal cortex and striatum. [1]
In vivo	The inhibitory actions of Lurasidone on MAP-induced hyperactivity persists for more than 8 hours, and the ED ₅₀ values of the action at 1 hour, 2 hours, 4 hours, and 8 hours after the treatment are 2.3 mg/kg, 0.87 mg/kg, 1.6 mg/kg, and 5.0 mg/kg, respectively. Lurasidone (1 mg/kg-10 mg/kg) dose-dependently inhibits conditioned avoidance response in rats with ED ₅₀ of 6.3 mg/kg. Lurasidone dose-dependently inhibits TRY-induced forepaw clonic seizure and p-CAMP-induced hyperthermia in rats with ED ₅₀ of 5.6 mg/kg and 3.0 mg/kg, respectively. Lurasidone (0.3 mg/kg-30 mg/kg) dose-dependently and significantly increases the number of shocks received by rats in the Vogel's conflict test with MED of 10 mg/kg. Lurasidone (3 mg/kg, 2 weeks) significantly suppresses hyperactivity behavior in olfactory bulbectomy model rats. Lurasidone (700 mg/kg-1000 mg/kg) slightly prolongs the duration of loss of righting reflexes elicited by hexobarbital (anesthesia) in mice in a dose-dependent manner. [1] Lurasidone (30 mg/kg, p.o.) significantly and dose-dependently reverses the MK-801-induced impairment of the passive-avoidance response of rats. [2] Lurasidone (3 mg/kg p.o.) potently reverses MK-801-induced learning impairment in the Morris water maze test in rats. Lurasidone (3 mg/kg p.o.) potently reverses MK-801-induced reference memory impairment and moderately but not significantly attenuates MK-801-induced working memory impairment in the radial-arm maze test. [3] Lurasidone (10 mg/kg) treatment increases total BDNF mRNA levels in rat prefrontal cortex and, to less extent, in hippocampus. Lurasidone (10 mg/kg) significantly increases the levels of mature BDNF

In vivo	protein in rat prefrontal cortex, without affect the protein levels of the neurotrophin (both precursor and mature forms) in hippocampal extracts. [4]
---------	--

Solubility Information

Solubility	DMSO: 6.96 mg/mL (13.15 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	---

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.8899 mL	9.4493 mL	18.8986 mL
5 mM	0.378 mL	1.8899 mL	3.7797 mL
10 mM	0.189 mL	0.9449 mL	1.8899 mL
50 mM	0.0378 mL	0.189 mL	0.378 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Ishibashi T, et al. *J Pharmacol Exp Ther*, 2010, 334(1), 171-181.

Danek P J, Daniel W A. The Atypical Antipsychotic Lurasidone Affects Brain but Not Liver Cytochrome P450 2D (CYP2D) Activity. A Comparison with Other Novel Neuroleptics and Significance for Drug Treatment of Schizophrenia. *Cells*. 2022, 11(21): 3513.

Ishiyama T, et al. *Eur J Pharmacol*, 2007, 572(2-3), 160-170.

Kot M, Haduch A, Papp M, et al. The effect of chronic treatment with lurasidone on rat liver cytochrome P450 expression and activity in the chronic mild stress (CMS) model of depression. *Drug Metabolism and Disposition*. 2017, 45(12): 1336-1344

Enomoto T, et al. *Behav Brain Res*, 2008, 186(2), 197-207.

Danek P J, Wójcikowski J, Daniel W A. The atypical neuroleptics iloperidone and lurasidone inhibit human cytochrome P450 enzymes in vitro. Evaluation of potential metabolic interactions. *Pharmacological Reports:PR*. 2020, 72(6): 1685-1694

Danek P J, Daniel W A. The effect of new atypical antipsychotic drugs on the expression of transcription factors regulating cytochrome P450 enzymes in rat liver. *Pharmacological Reports*. 2024: 1-7.

Fumagalli F, et al. *Int J Neuropsychopharmacol*, 2012, 15(2), 235-246.

Danek P J, Wójcikowski J, Daniel W A. The atypical neuroleptics iloperidone and lurasidone inhibit human cytochrome P450 enzymes in vitro. Evaluation of potential metabolic interactions[J]. *Pharmacological reports: PR*. 2020, 72(6): 1685-1694.

Marta Kot, Anna Haduch, Mariusz Papp, Władysława A. Daniel. The effect of chronic treatment with lurasidone on rat liver cytochrome P450 expression and activity in the chronic mild stress (CMS) model of depression [J]. *Drug Metabolism and Disposition*. 2017, 45(12): 1336-1344.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481