

## AMI-193

## Chemical Properties


CAS No. : 510-74-7

Formula: C<sub>22</sub>H<sub>26</sub>N<sub>3</sub>O<sub>2</sub>

Molecular Weight: 383.46

Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year

Actual storage temperature shall be subject to the COA.



## Biological Description

Description	AMI-193 (Spiramide) is an effective and selective antagonist of 5-HT <sub>2</sub> and D <sub>2</sub> receptor with K <sub>i</sub> s of 2, 3, 50, 2530, and 4300 nM for 5-HT <sub>2</sub> , D <sub>2</sub> Receptor, 5-HT <sub>1A</sub> , D <sub>1</sub> Receptor, and 5-HT <sub>1C</sub> Receptor. AMI-193 shows antipsychotic activity.
Targets(IC <sub>50</sub> )	5-HT Receptor, Dopamine Receptor
In vivo	In adult male squirrel monkeys, AMI-193 (0.003 and 0.01 mg/kg; i.m.) decreases the response rate under a fixed-interval (FI) schedule of stimulus termination in a dose-dependent manner and under a second-order schedule of i.v. self-administration of cocaine (0.1 mg/infusion). AMI-193 (0.003 and 0.01 mg/kg; i.m.) attenuates the discriminative-stimulus effects of cocaine[2].

## Solubility Information

Solubility	DMSO: 20.7 mg/mL (53.98 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 2 mg/mL (5.22 mM), Sonication is recommended. 10% DMSO+90% Saline: 2.07 mg/mL (5.4 mM), Solution. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

### Preparing Stock Solutions

---

	1mg	5mg	10mg
1 mM	2.6078 mL	13.0392 mL	26.0783 mL
5 mM	0.5216 mL	2.6078 mL	5.2157 mL
10 mM	0.2608 mL	1.3039 mL	2.6078 mL
50 mM	0.0522 mL	0.2608 mL	0.5216 mL

---

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

### Reference

- Kjellberg B, et, al. Partial restoration by a neuroleptic (spiramide) of items of grooming behaviour suppressed by amphetamine. *Arch Int Pharmacodyn Ther.* 1974 Jul;210(1):61-6.
- Ismail AM, et, al. Antagonism of 1-(2,5-dimethoxy-4-methylphenyl)-2-aminopropane stimulus with a newly identified 5-HT<sub>2</sub>- versus 5-HT<sub>1C</sub>-selective antagonist. *J Med Chem.* 1993 Aug 20;36(17):2519-25.
- Czoty PW, et, al. Behavioral effects of AMI-193, a 5-HT(2A)- and dopamine D(2)-receptor antagonist, in the squirrel monkey. *Pharmacol Biochem Behav.* 2000 Oct;67(2):257-64.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481