

BOC-NH-PEG2-propene

Chemical Properties

CAS No. : 2410236-85-8


Formula: C₁₂H₂₃N₁O₄

Molecular Weight: 245.319

Keep away from direct sunlight

Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year

Actual storage temperature shall be subject to the COA.



Biological Description

Description	BOC-NH-PEG2-propene is a PEG-based linker used in PROTACs to join two essential ligands, facilitating the formation of PROTAC molecules. This linker enables selective protein degradation by leveraging the ubiquitin-proteasome system within cells.
Targets(IC50)	Others,PROTAC Linker
In vitro	PROTACs consist of two distinct ligands linked together; one targets an E3 ubiquitin ligase and the other binds to the target protein. By leveraging the intracellular ubiquitin-proteasome system, PROTACs selectively induce the degradation of target

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	4.0763 mL	20.3815 mL	40.7631 mL
5 mM	0.8153 mL	4.0763 mL	8.1526 mL
10 mM	0.4076 mL	2.0382 mL	4.0763 mL
50 mM	0.0815 mL	0.4076 mL	0.8153 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

An S, et al. Small-molecule PROTACs: An emerging and promising approach for the development of targeted therapy drugs. EBioMedicine. 2018 Oct;36:553-562

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481