


D,L-erythro-PDMP

Chemical Properties

CAS No. :	109760-77-2
Formula:	C ₂₃ H ₃₈ N ₂ O ₃
Molecular Weight:	390.568
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



Biological Description

Description	D,L-erythro-PDMP, an erythro isomer of PDMP, exhibits growth inhibition on cultured rabbit skin fibroblasts. Additionally, this compound serves as an effective inhibitor of UDP-glucose: ceramide glucosyltransferase.
Targets(IC50)	Others, Transferase
In vitro	D,L-erythro-PDMP at concentrations of 0, 12, 25, 50 μ M over periods of 0, 4, 7, and 10 days demonstrates a dose-dependent inhibition of rabbit skin fibroblast proliferation, as evidenced by reduced cell growth[1]. A concentration of 50 μ M of D,L-erythro-PDMP, when applied for 3 days, results in cytotoxic effects, altering the cell morphology of rabbit skin fibroblasts[1]. Furthermore, treatment with D,L-erythro-PDMP at a concentration of 40 μ M for 24 hours in MDCK cells leads to a significant increase in glucosyltransferase activity, quantified at 14.6 nmol/h per mg of protein[2]. Additionally, a 40 μ M concentration of D,L-erythro-PDMP administered for 6 hours offers protective effects against synthase loss in cells challenged with cycloheximide[2]. These findings are supported by cell proliferation and cytotoxicity assays on rabbit skin fibroblast cell lines, illustrating D,L-erythro-PDMP's capacity to modulate cell growth and viability in a concentration and time-dependent manner[1].

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.5604 mL	12.8018 mL	25.6036 mL
5 mM	0.5121 mL	2.5604 mL	5.1207 mL
10 mM	0.256 mL	1.2802 mL	2.5604 mL
50 mM	0.0512 mL	0.256 mL	0.5121 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Uemura K, et al. Effect of an inhibitor of glucosylceramide synthesis on cultured rabbit skin fibroblasts. J Biochem. 1990 Oct;108(4):525-30.

Abe A, et al. Induction of glucosylceramide synthase by synthase inhibitors and ceramide. Biochim Biophys Acta. 1996 Feb 16;1299(3):333-41.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481