Data Sheet (Cat.No.T1724)



SCR7 pyrazine

Chemical Propert	ies
CAS No. :	14892-97-8
Formula:	C18H12N4OS
Molecular Weight:	332.38
Appearance:	no data available
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year

Biological Description

SCR7 pyrazine (SCR7) enhances CRISPR-Cas9-mediated homology-directed repair (H efficiency in vitro up to 19-fold. Inhibits nonhomologous end-joining (NHEJ).		
Apoptosis,DNA/RNA Synthesis,CRISPR/Cas9		
SCR7 effectively inhibits the formation of multimers at 200 µM and above. SCR7 successfully inhibits cell proliferation of MCF7, A549, HeLa, T47D, A2780, HT1080, and Nalm6 with IC50 of 40, 34, 44, 8.5, 120, 10, and 50 µM, respectively.[1] SCR7 suppresses the NHEJ repair of CRISPR-Cas9-induced DSBs.[2]		
SCR7 treatment (10 mg/kg, i.m.) signi?cantly reduces breast adenocarcinoma-induced tumor, and exhibits 4-fold increase in lifespan compared with control group. However, in Swiss albino mice with Dalton's lymphoma tumor model, SCR7 (20 mg/kg, i.p.) exhibits neither tumor regression nor increase in lifespan. In BALB/c mice, SCR7 (20 mg/kg, i.p.) signi?cantly enhances the cytotoxic effects of radiation, etoposide and 3- Aminobenzamide on tumor derived from Dalton's lymphoma (DLA) cells.[1]		
Complementation of SCR7 Inhibition with Puri?ed Ligase IV: Complementation experiment is carried out by adding increasing concentrations of puri?ed Ligase IV/XRCC4 complex (30, 60, and 120 fmol) along with the oligomeric DNA substrates (5' compatible and 5'-5' noncompatible ends) to the SCR7-treatedextracts. Reactions are incubated for 2 h at 25°C. The reaction products are then resolved on 8% denaturing PAGE. The gel is dried and exposed and the signal is detected with a PhosphorImager and analyzed with Multi Gauge (V3.0) software.		
Cell proliferation of cancer cells are determined by MTT and trypan blue assays. Briefly, MCF7, CEM, HeLa, A549, HT1080, A2780, T47D, Nalm6, N114 and K562 cells are grown in presence of SCR7 (10, 50, 100, and 250?µM) for 24 or 48?h, and subjected to MTT or trypan blue assays. Each experiment is repeated a minimum of three independent times. (Only for Reference)		

Solubility Information	
Solubility	H2O: <1 mg/mL,
	Ethanol: 3 mg/mL (9.03 mM),
	DMSO: 66 mg/mL (198.57 mM),
	(< 1 mg/ml refers to the product slightly soluble or insoluble)

A DRUG SCREENING EXPERT

Preparing Stock Solutions

	1mg	5mg	10mg	
1 mM	3.0086 mL	15.043 mL	30.086 mL	
5 mM	0.6017 mL	3.0086 mL	6.0172 mL	
10 mM	0.3009 mL	1.5043 mL	3.0086 mL	
50 mM	0.0602 mL	0.3009 mL	0.6017 mL	

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Srivastava M, et al. Cell. 2012, 151(7), 1474-1487. Chu VT, et al. Nat Biotechnol. 2015, doi: 10.1038/nbt.3198.

Inhibitor • Natural Compounds • Compound Libraries • Recombinant Proteins This product is for Research Use Only• Not for Human or Veterinary or Therapeutic Use Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481