


Direct Black 38 free acid

Chemical Properties

CAS No. :	22244-14-0
Formula:	C34H27N9O7S2
Molecular Weight:	737.77
Appearance:	no data available
Storage:	keep away from direct sunlight Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Ferristatin II (Direct Black 38 free acid) is a polysulphonated dye and promotes the degradation of transferrin receptor-1 in vitro and in vivo.
Targets(IC50)	Others

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.3554 mL	6.7772 mL	13.5544 mL
5 mM	0.2711 mL	1.3554 mL	2.7109 mL
10 mM	0.1355 mL	0.6777 mL	1.3554 mL
50 mM	0.0271 mL	0.1355 mL	0.2711 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Beije B. Induction of unscheduled DNA synthesis in liver and micronucleus in bone marrow of rats exposed in vivo to the benzidine-derived azo dye, Direct Black 38. *Mutat Res.* 1987 Apr;187(4):227-34.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481