


Taselisib

Chemical Properties

CAS No. :	1282512-48-4
Formula:	C ₂₄ H ₂₈ N ₈ O ₂
Molecular Weight:	460.53
Appearance:	no data available
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Taselisib (GDC-0032) is an orally bioavailable inhibitor of the class I phosphatidylinositol 3-kinase (PI3K) alpha isoform (PIK3CA), with potential antineoplastic activity.
Targets(IC50)	PI3K,Carbonic Anhydrase
Kinase Assay	Characterization of Biochemical and Cellular Activity in Vitro: Enzymatic activity of the class I PI3K isoforms is measured using a fluorescence polarization assay that monitors formation of the product 3,4,5-inositoltriphosphate molecule as it competes with fluorescently labeled PIP3 for binding to the GRP-1 pleckstrin homology domain protein. An increase in phosphatidyl inositide-3-phosphate product results in a decrease in fluorescence polarization signal as the labeled fluorophore is displaced from the GRP-1 protein binding site. Class I PI3K isoforms are expressed and purified as heterodimeric recombinant proteins. Tetramethylrhodamine-labeled PIP3 (TAMRA-PIP3), di-C8-PIP2, and PIP3 detection reagents are purchased from Echelon Biosciences. PI3K α is assayed under initial rate conditions in the presence of 10 mM Tris (pH 7.5), 25 μ M ATP, 9.75 μ M PIP2, 5% glycerol, 4 mM MgCl ₂ , 50 mM NaCl, 0.05% (v/v) Chaps, 1 mM dithiothreitol, and 2% (v/v) DMSO at 60 ng/mL. After assay for 30 min at 25 °C, reactions are terminated with a final concentration of 9 mM EDTA, 4.5 nM TAMRA-PIP3, and 4.2 μ g/mL GRP-1 detector protein before reading fluorescence polarization on an Envision plate reader. IC ₅₀ values are calculated from the fit of the dose-response curves to a 4-parameter equation. Each reported value is an average of three experiments, and all have a standard deviation less than one geometric mean.
Cell Research	GDC-0032 is dissolved in DMSO. Cells are seeded in replicates of 6 in 96-well plates with 500 to 5,000 cells/well overnight and then treated with GDC-0032. After 4 days, the media are removed and the cells are fixed with 4% glutaraldehyde for 30 minutes. Fixed cells are stained with 0.1% crystal violet for 2 minutes, then washed, and dissolved in 10% acetic acid.

Solubility Information

Solubility	DMSO: 70 mg/mL (151.99 mM),Heating is recommended. Ethanol: < 1 mg/mL (insoluble or slightly soluble), < 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.1714 mL	10.8571 mL	21.7141 mL
5 mM	0.4343 mL	2.1714 mL	4.3428 mL
10 mM	0.2171 mL	1.0857 mL	2.1714 mL
50 mM	0.0434 mL	0.2171 mL	0.4343 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Huang Q, Ru Y, Luo Y, et al. Identification of a targeted ACSL4 inhibitor to treat ferroptosis-related diseases. Science Advances. 2024, 10(13): eadk1200.