


Menin-MLL inhibitor MI-2

Chemical Properties

CAS No. :	1271738-62-5
Formula:	C ₁₈ H ₂₅ N ₅ S ₂
Molecular Weight:	375.55
Appearance:	no data available
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Menin-MLL inhibitor MI-2 (MI2) is a potent menin-MLL interaction inhibitor with IC ₅₀ of 446 nM.
Targets(IC ₅₀)	Apoptosis,Epigenetic Reader Domain,Histone Methyltransferase
In vivo	MI-2 effectively inhibits cell proliferation and induces apoptosis in human MLL leukemia cell lines bearing various MLL translocations. In HEK293 cells, MI-2 penetrates its protein target, efficiently inhibiting the interaction between menin and MLL-AF9. By downregulating the expression of target genes essential for the oncogenic activity of MLL fusion proteins, MI-2 effectively blocks leukemia transformation mediated by MLL fusion proteins.
Kinase Assay	High Throughput Screening: FITC-MBM1 at 15 nM and menin at 150 nM in the FP buffer are mixed and incubated for 1h in the dark at room temperature. For point screening, the 0.2 µL of each compound (20 µM final concentration, 1% DMSO) is added to 20 µL of the aliquot of the protein-peptide mixture and incubated on 384-well plates in the dark at room temperature for 1h. In confirmation screening, the serial dilution plates with compounds in DMSO are prepared and used to titrate the menin-FITC-MBM1 complex. Change in fluorescence polarization is monitored at 525 nm after excitations at 495 nm using the PHERAstar microplate reader (BMG) and applied to determine IC ₅₀ values with the Origin 7.0 program.
Cell Research	The MLL-AF9 and E2A-HLF transduced murine BMC are plated in 12-well plates at the concentration of 5×10 ³ cells/mL with 1 mL methylcellulose medium M3234 containing 20% IMDM medium, 1% penicillin/streptomycin, IL-3 and 0.25% DMSO or compounds. 6 days later colonies are stained with 100 µL iodonitrotetrazolium chloride at final concentration of 1 mg/mL, incubated at 37°C for 30 min and counted. To replat for the 2nd round, colonies are counted at day 6 without staining and cells were washed out by 1×PBS buffer and resuspended in IMDM medium containing 15% FBS, 1% penicillin/streptomycin and IL-3. 5×10 ³ cells are plated in 12-well plates with 1ml methylcellulose medium M3234 containing 20% IMDM medium, 1% penicillin/streptomycin, IL-3 and 0.25% DMSO or compounds. 6 days later colonies are stained and counted.(Only for Reference)

Solubility Information

A DRUG SCREENING EXPERT

Solubility	DMSO: 70 mg/mL (186.4 mM), H2O: < 1 mg/mL (insoluble or slightly soluble), Ethanol: 70 mg/mL (186.4 mM), < 1 mg/ml refers to the product slightly soluble or insoluble)
------------	---

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.6628 mL	13.3138 mL	26.6276 mL
5 mM	0.5326 mL	2.6628 mL	5.3255 mL
10 mM	0.2663 mL	1.3314 mL	2.6628 mL
50 mM	0.0533 mL	0.2663 mL	0.5326 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Grembecka J, et al. Nat Chem Biol. 2012, 8(3), 277-284.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481