

Lipoteichoic acid

Chemical Properties

CAS No. : 56411-57-5

Formula:

Molecular Weight:

Appearance: no data available

Storage: keep away from direct sunlight
Powder: -20°C for 3 years | In solvent: -80°C for 1 year

Lipoteichoic acid

Biological Description

Description	Lipoteichoic acid, a polymer of Staphylococcus aureus cell wall, activates the complement system by inducing C3 and inhibiting CD55, and reduces fat deposition via the IGF-1 pathway.
Targets(IC50)	Complement System

Solubility Information

Solubility	DMSO: 80 mg/mL, Sonication is recommended. H2O: 5 mg/mL, Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	---

Reference

Bong Jun Jung, et al. Lipoteichoic Acid from Staphylococcus aureus Activates the Complement System via C3 Induction and CD55 Inhibition. Microorganisms. 2021 May 24;9(6):1135.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 36 Washington Street, Wellesley Hills, MA 02481