


Phalloidin-TRITC

Chemical Properties

CAS No. :	915013-10-4
Formula:	C60H70N12O13S2
Molecular Weight:	1231.41
Appearance:	no data available
Storage:	keep away from direct sunlight Powder: -20°C for 3 years In solvent: -80°C for 1 year



Biological Description

Description	Phalloidin-TRITC is a TRITC labeled, red fluorescence probe for F-actin . Phalloidin, bound to actin filaments, reacts covalently with amino acids Glu-IIT, Met-II9, and Met355, which are very close to the nucleotide binding site.
In vitro	Phalloidin induced actin polymerization in the cytoplasm of cultured cells interferes with cell locomotion and growth[2].

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	0.8121 mL	4.0604 mL	8.1208 mL
5 mM	0.1624 mL	0.8121 mL	1.6242 mL
10 mM	0.0812 mL	0.406 mL	0.8121 mL
50 mM	0.0162 mL	0.0812 mL	0.1624 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

- J A Cooper, et al. Effects of cytochalasin and phalloidin on actin. J Cell Biol. 1987 Oct;105(4):1473-8.
- J Wehland, et al. Phalloidin-induced actin polymerization in the cytoplasm of cultured cells interferes with cell

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481