


UK-5099

Chemical Properties

CAS No. : 56396-35-1
Formula: C₁₈H₁₂N₂O₂
Molecular Weight: 288.3
Appearance: no data available
Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year



Biological Description

Description	UK-5099 (PF-1005023) is a specific and potent inhibitor of MPC carrier activity [1]. It potently inhibits pyruvate-dependent oxygen consumption by rat heart mitochondria (IC ₅₀ : 50 nM) [1].
Targets(IC ₅₀)	Others,Mitochondrial Metabolism
In vitro	The trypanosomal pyruvate carrier is found to be rather insensitive to inhibition by alpha-cyano-4-hydroxycinnamate (K _i : 17 mM) but can be completely blocked by UK-5099 (K _i : 49 μM)[2]. UK-5099 also inhibits the monocarboxylate transporter (MCT) [3]. UK5099 dose-dependently significantly inhibits the glucose-stimulated rise in oxygen consumption and at 150 μM reduced oxygen consumption below basal levels. UK5099 reduces ATP levels and increases ADP and AMP levels in 832/13 cells[4]. The UK5099 treated cells show a significantly higher proportion of side population fraction and express higher levels of stemness markers Oct3/4 and Nanog. The UK5099 application may be an ideal model for Warburg effect studies[5].
In vivo	UK5099 increases the glucose excursion seen during an intraperitoneal glucose tolerance test in C57BLK mice[4].
Cell Research	The 832/13 cell line derived from INS-1 rat insulinoma cells is used for experiments. Cell viability is measured using CellTiter Blue. The assay is based on the cellular reduction of resazurin to resorufin. The appearance of resorufin is monitored by fluorescence emission at 585 nm using a Spectramax M5 microplate reader with excitation at 555 nm. For UK5099-treated cells, cells are allowed to recover for 1 h before measuring cell viability. Data are expressed as -fold relative to no treatment or siCtrl[4].
Animal Research	C57BLK mice are fasted for 16 h prior to glucose challenge. UK5099 (32 μmol/kg of body weight) or DMSO in PBS is injected into the intraperitoneal cavity 30 min before injecting glucose (1.5 mg of glucose/g of body weight). Blood glucose levels are measured at 0, 10, 20, 30, 60, and 120 min after glucose injection[4].

Solubility Information

Solubility	DMSO: 16.67 mg/mL (57.81 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.4686 mL	17.343 mL	34.6861 mL
5 mM	0.6937 mL	3.4686 mL	6.9372 mL
10 mM	0.3469 mL	1.7343 mL	3.4686 mL
50 mM	0.0694 mL	0.3469 mL	0.6937 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Halestrap A P. The mitochondrial pyruvate carrier. Kinetics and specificity for substrates and inhibitors[J]. Biochemical Journal, 1975, 148(1): 85-96.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481