Data Sheet (Cat.No.T6452)



Closantel Sodium

Chemical Properties


CAS No.: 61438-64-0

Formula: C22H13Cl2I2N2O2·Na

Molecular Weight: 685.06

Appearance: no data available

Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year



Biological Description

| Description | Closantel is a salicylanilide anthelmintic compound; exhibits different anthelmintic spectra and apparent toxicity in mammals. |
|---------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Targets(IC50) | Antibacterial, Parasite |
| In vitro | Closantel inhibits drug resistant S. aureus and E. faecium with MICs of 1 µg/mL and 1 µg/mL. Closantel inhibits MRSA and E.faecalis with MICs of 2 µg/mL and 1 µg/mL. [1] Closantel completely blocks the infection of host cells with E. chaffeensis or A. phagocytophilum, treatment of infected cells 1 day post infection with closantel clears infection in dose-dependent manner. Closantel inhibits the autokinase activities of the three E. chaffeensis sensor kinases from E. chaffeensis. [2] Closantel (50 µg/mL) causes an initial burst of contractions of much greater amplitude and frequency than normal in tissue vessel, the stimulation of amplitude and frequency lasted for nearly 15 min and is accompanied by a rise in muscle tone, which reaches a maximum at 16 min, at a level greater than 1.5 times the maximal normal amplitude. [3] |
| In vivo | Closantel (10 mg/kg) results in gross surface damage from 24 hours onwards in rats administrated with 25 metacercarial cysts of F. hepatica, in the form of erosion of the anterior and posterior extremities of the fluke and large-scale sloughing of the tegument on both dorsal and ventral surfaces. [3] Closantel (7.5 mg/kg) combined with broad-spectrum anthelmintic is very effective against H. contortus but ineffective against Trichostrongylus spp in lambs. [4] Closantel significantly reduces isotope levels in closantel-resistant adult Haemonchus contortus infected sheep. [5] |

Solubility Information

| Solubility | H2O: < 1 mg/mL (insoluble or slightly soluble), Omso: 93 mg/mL (135.8 |
|------------|------------------------------------------------------------------------------|
| | mM), Ethanol: 93 mg/mL (135.8 mM), (< 1 mg/ml refers to the product |
| | slightly soluble or insoluble) |

Page 1 of 2 www.targetmol.com

Preparing Stock Solutions

| | 1mg | 5mg | 10mg |
|-------|-----------|-----------|------------|
| 1 mM | 1.4597 mL | 7.2986 mL | 14.5973 mL |
| 5 mM | 0.2919 mL | 1.4597 mL | 2.9195 mL |
| 10 mM | 0.146 mL | 0.7299 mL | 1.4597 mL |
| 50 mM | 0.0292 mL | 0.146 mL | 0.2919 mL |

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Hlasta DJ, et al. Bioorg Med Chem Lett, 1998, 8(14), 1923-1928. Cheng Z, et al. Cell Microbiol, 2006, 8(8), 1241-1252.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481

Page 2 of 2 www.targetmol.com