Data Sheet (Cat.No.T6575)



LY2119620

Chemical Properties

CAS No.: 886047-22-9

Formula: C19H24ClN5O3S

Molecular Weight: 437.94

Appearance: no data available

Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year

acetylcholine receptors.

$\bigcup_{N \nmid 1} \bigcup_{H,N} S \bigvee_{CH_1} \bigcup_{CH_2} \bigcup_{CH_3} \bigcup_{CH_4} \bigcup_{CH_3} \bigcup_{CH_4} \bigcup_{CH_4} \bigcup_{CH_3} \bigcup_{CH_4} \bigcup_{CH_4} \bigcup_{CH_4} \bigcup_{CH_4} \bigcup_{CH_4} \bigcup_{CH_5} \bigcup_{CH_5}$

Biological Description

Description

M2 and M4 receptors into an active G protein-bound state. LY2119620 enhances the potency of three muscarinic acetylcholine receptor agonists, ACh, Oxo-M and iperoxo. [1] [3H]LY2119620 can be used as a probe for the human M(2) and M(4) muscarinic receptor allosteric binding sites. [2] Kinase Assay Biochemical Assay: For IC50 determination, EI1 is serial diluted threefold in DMSO for a total of 12 concentrations, with the starting concentration at 1 μM. The reaction is incubated at room temperature for 120 min, and stopped by adding quench solution (2.5% TFA with 320 nM d4-SAH). SAH production is quantitated using an API 4000 triple quadrupole mass spectrometry with Turbulon Spray coupled with Prominence UFLC. The percentage of inhibition is normalized using positive (no inhibitor) and negative (no enzyme) controls, and IC50 calculated using PRISM. Enzymology studies of S-Adenosyl methionine (SAM) competition are performed with slight modification of reaction condition: 10 μM EI1 is used as the starting dose for serial dilution. SAM is titrated over a range between 1 μM and 50 μM (corresponding to 1 × Km and 50 × Km), and substrate peptide is present in the final reaction mixture at its saturated condition (10 μM). For histone methyltransferase (HMT) profiling in Table 1, all HMTs are purified recombinant		dectylenoune receptors.
G protein-signaling ability of an agonist at the M2/M4 receptor subtypes by placing the M2 and M4 receptors into an active G protein-bound state. LY2119620 enhances the potency of three muscarinic acetylcholine receptor agonists, ACh, Oxo-M and iperoxo. [1] [3H]LY2119620 can be used as a probe for the human M(2) and M(4) muscarinic receptor allosteric binding sites. [2] Kinase Assay Biochemical Assay: For IC50 determination, EI1 is serial diluted threefold in DMSO for a total of 12 concentrations, with the starting concentration at 1 μM. The reaction is incubated at room temperature for 120 min, and stopped by adding quench solution (2.5% TFA with 320 nM d4-SAH). SAH production is quantitated using an API 4000 triple quadrupole mass spectrometry with Turbulon Spray coupled with Prominence UFLC. The percentage of inhibition is normalized using positive (no inhibitor) and negative (no enzyme) controls, and IC50 calculated using PRISM. Enzymology studies of S-Adenosyl methionine (SAM) competition are performed with slight modification of reaction condition: 10 μM EI1 is used as the starting dose for serial dilution. SAM is titrated over a range between 1 μM and 50 μM (corresponding to 1 × Km and 50 × Km), and substrate peptide is present in the final reaction mixture at its saturated condition (10 μM). For histone methyltransferase (HMT) profiling in Table 1, all HMTs are purified recombinant proteins from either Escherichia coli or Baculovirus system. The catalytic domain of G9a SuV39H2, Set7/9, CARM1, SETD8, NSD3, SETD2, and Dot1L, and the full-length SmyD2 protein were used in the biochemical assays. HMT biochemical reactions are carefully	Targets(IC50)	AChR
total of 12 concentrations, with the starting concentration at 1 µM. The reaction is incubated at room temperature for 120 min, and stopped by adding quench solution (2.5% TFA with 320 nM d4-SAH). SAH production is quantitated using an API 4000 triple quadrupole mass spectrometry with Turbulon Spray coupled with Prominence UFLC. The percentage of inhibition is normalized using positive (no inhibitor) and negative (no enzyme) controls, and IC50 calculated using PRISM. Enzymology studies of S-Adenosyl methionine (SAM) competition are performed with slight modification of reaction condition: 10 µM EI1 is used as the starting dose for serial dilution. SAM is titrated over a range between 1 µM and 50 µM (corresponding to 1 × Km and 50 × Km), and substrate peptide is present in the final reaction mixture at its saturated condition (10 µM). For histone methyltransferase (HMT) profiling in Table 1, all HMTs are purified recombinant proteins from either Escherichia coli or Baculovirus system. The catalytic domain of G9a SuV39H2, Set7/9, CARM1, SETD8, NSD3, SETD2, and Dot1L, and the full-length SmyD2 protein were used in the biochemical assays. HMT biochemical reactions are carefully		G protein-signaling ability of an agonist at the M2/M4 receptor subtypes by placing the M2 and M4 receptors into an active G protein-bound state. LY2119620 enhances the potency of three muscarinic acetylcholine receptor agonists, ACh, Oxo-M and iperoxo. [1] [3H]LY2119620 can be used as a probe for the human M(2) and M(4) muscarinic
i characienzeu wini enzymbiouv studies and the part and substidie XIII determined. The	Kinase Assay	total of 12 concentrations, with the starting concentration at 1 μ M. The reaction is incubated at room temperature for 120 min, and stopped by adding quench solution (2.5% TFA with 320 nM d4-SAH). SAH production is quantitated using an API 4000 triple quadrupole mass spectrometry with Turbulon Spray coupled with Prominence UFLC. The percentage of inhibition is normalized using positive (no inhibitor) and negative (no enzyme) controls, and IC50 calculated using PRISM. Enzymology studies of S-Adenosyl methionine (SAM) competition are performed with slight modification of reaction condition: 10 μ M EI1 is used as the starting dose for serial dilution. SAM is titrated over a range between 1 μ M and 50 μ M (corresponding to 1 × Km and 50 × Km), and substrate peptide is present in the final reaction mixture at its saturated condition (10 μ M). For histone methyltransferase (HMT) profiling in Table 1, all HMTs are purified recombinant proteins from either Escherichia coli or Baculovirus system. The catalytic domain of G9a, SuV39H2, Set7/9, CARM1, SETD8, NSD3, SETD2, and Dot1L, and the full-length SmyD2 protein were used in the biochemical assays. HMT biochemical reactions are carefully
SAM and substrate concentrations are kept at their respective Km for most of the HMTs, except the ones (SmyD2 and Set7/9) with low SAM-Km value, for which 0.5 µM SAM is		SAM and substrate concentrations are kept at their respective Km for most of the HMTs,
used. All HMT reactions are performed using the same assay format where the production of SAH from the biochemical reaction is quantitated by LC-MS.		used. All HMT reactions are performed using the same assay format where the

LY2119620 is a specific, and allosteric agonist of human M2 and M4 muscarinic

Solubility Information

A DRUG SCREENING EXPERT

Solubility	DMSO: 50 mg/mL (114.17 mM),	70
	Ethanol: 10 mg/mL, Heating is recommended.	
	(< 1 mg/ml refers to the product slightly soluble or insoluble)	

Preparing Stock Solutions

	1mg	5mg	10mg	
1 mM	2.2834 mL	11.4171 mL	22.8342 mL	
5 mM	0.4567 mL	2.2834 mL	4.5668 mL	
10 mM	0.2283 mL	1.1417 mL	2.2834 mL	
50 mM	0.0457 mL	0.2283 mL	0.4567 mL	

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Croy CH, et al. Mol Pharmacol. 2014, 86(1), 106-115. Schober DA, et al. Mol Pharmacol. 2014, 86(1), 116-123.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481

Page 2 of 2 www.targetmol.com