Data Sheet (Cat.No.T652023)



Harpagoside

Chemical Proper	ties	
CAS No. :	19210-12-9	HO
Formula:	C24H30O11	но
Molecular Weight:	494.49	
Appearance: 🦲	no data available	Н ₃ С ОН
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year	но

Biological Description	
Description	1. Harpagoside has anti-inflammatory activity. 2. Harpagoside at 3 mM concentration, shows moderate inhibition of seed germination. 3. Harpagoside can inhibit lipopolysaccharide-induced mRNA levels and protein expression of cyclooxygenase-2 and inducible nitric oxide in HepG2 cells.
Targets(IC50)	Others,COX,NO Synthase

Solubility Information	
Solubility	Ethanol: Soluble, DMSO: 90 mg/mL(182 mM), Pyridine, Methanol, etc.: Soluble, (< 1 mg/ml refers to the product slightly soluble or insoluble)

Preparing Stock Solutions

<u>e</u>	1mg	5mg	10mg
1 mM	2.0223 mL	10.1114 mL	20.2229 mL
5 mM	0.4045 mL	2.0223 mL	4.0446 mL
10 mM	0.2022 mL	1.0111 mL	2.0223 mL
50 mM	0.0404 mL	0.2022 mL	0.4045 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Kim T K , Park K S . Inhibitory effects of harpagoside on TNF-α-induced pro-inflammatory adipokine expression through PPAR-γ activation in 3T3-L1 adipocytes[J]. Cytokine, 2015, 76(2):S104346661500201X.

Inhibitor • Natural Compounds • Compound Libraries • Recombinant ProteinsThis product is for Research Use Only• Not for Human or Veterinary or Therapeutic UseTel:781-999-4286E_mail:info@targetmol.comAddress:36 Washington Street,Wellesley Hills,MA 02481