

SARS-CoV-2 Spike S1+S2 Protein (157-158 deletion & T19R & G142D & E156G & L452R & T478K & H156R & N501Y & P681R), Biotinylated

General Information

Protein Construction: The SARS-CoV-2 Spike S1+S2 (YP_009724390.1, with mutations 157-158 deletion, T19R, G142D, E156G, L452R, T478K, D614G, P681R, D950N) (Met1-Pro1213) was expressed with a C-terminal polyhistidine tag followed by an AVI tag. The expressed protein was biotinylated in vivo by the Biotin-Protein ligase (BirA enzyme) which is co-expressed. The mutations were identified in the SARS-CoV-2 variant (known as B.1.617.2) which emerged in the India.

Species: SARS-CoV-2

Expression Host: Baculovirus Insect Cells

Accession: A0A6G7K2L4

Molecular Weight: 136.01 kDa (predicted)

QC Testing

Biological Activity: Testing in progress

Purity: > 90 % as determined by SDS-PAGE.

Endotoxin: < 1.0 EU/μg of the protein as determined by the LAL method.

Formulation: Lyophilized from a solution filtered through a 0.22 μm filter, containing 20 mM Tris, 300 mM NaCl, 10% glycerol, pH 8.0. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:
A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:
It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted proteinsolutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Shipping:
In general, Lyophilized powders are shipping with blue ice.

Protein Background

The spike (S) glycoprotein of coronaviruses contains protrusions that will only bind to certain receptors on the host cell. Known receptors bind S1 are ACE2, angiotensin-converting enzyme 2; DPP4, dipeptidyl peptidase-4; APN, aminopeptidase N; CEACAM, carcinoembryonic antigen-related cell adhesion molecule 1; Sia, sialic acid; O-ac Sia, O-acetylated sialic acid. The spike is essential for both host specificity and viral infectivity. The term 'peplomer' is typically used to refer to a grouping of heterologous proteins on the virus surface that function together. The

spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection process. It's been reported that SARS-CoV-2 (COVID-19 coronavirus, 2019-nCoV) can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity. The main functions for the Spike protein are summarized as: Mediate receptor binding and membrane fusion; Defines the range of the hosts and specificity of the virus; Main component to bind with the neutralizing antibody; Key target for vaccine design; Can be transmitted between different hosts through gene recombination or mutation of the receptor binding domain (RBD), leading to a higher mortality rate.

Reference

Xiao X,et al. (2004) The SARS-CoV S glycoprotein. Cell Mol Life Sci. 61 (19-20): 2428-30.

Shen S,et al. (2007) Expression, glycosylation, and modification of the spike (S) glycoprotein of SARS CoV. Methods

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481