Data Sheet (Cat.No.TN4583)



Moringin

Chemical Proper	ties
CAS No. :	73255-40-0
Formula:	C14H17N05S
Molecular Weight:	311.35
Appearance:	no data available
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year

Biological Description

Description	Moringin is an isothiocyanate obtained from myrosinase-mediated hydrolysis of the glucosinolate precursor glucoMoringin present in Moringa oleifera seeds. Moringin has anti-inflammatory and antioxidative properties.
Targets(IC50)	gp120/CD4,IL Receptor,BCL,GSK-3,TNF,Wnt/beta-catenin,Caspase,COX,Nrf2,PPAR,p53
In vitro	 Moringin modulates neuroinflammation via the β-catenin-PPARγ axis and activates Wnt canonical pathway by inhibiting GSK3β in a mouse model of experimental autoimmune encephalomyelitis. Moringin is effective in inducing apoptosis through p53 and Bax activation and Bcl-2 inhibition. Moringin demonstrates the antitumor efficacy in human malignant astrocytoma cells. The combination of Cannabidiol (5µM) and Moringin (5µM) has anti-inflammatory, antioxidative, and anti-apoptotic effects[1].

Solubility Information

Solubility	DMSO: 62.5 mg/mL (200.74 mM),Sonication is recommended.	
	(< 1 mg/ml refers to the product slightly soluble or insoluble)	

Preparing Stock Solutions

	1mg	5mg	10mg	
1 mM	3.2118 mL	16.0591 mL	32.1182 mL	
5 mM	0.6424 mL	3.2118 mL	6.4236 mL	
10 mM	0.3212 mL	1.6059 mL	3.2118 mL	
50 mM	0.0642 mL	0.3212 mL	0.6424 mL	

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Rajan TS, et al. Anti-inflammatory and antioxidant effects of a combination of cannabidiol and moringin in LPSstimulated macrophages. Fitoterapia. 2016 Jul;112:104-15.

Inhibitor • Natural Compounds • Compound Libraries • Recombinant ProteinsThis product is for Research Use Only• Not for Human or Veterinary or Therapeutic UseTel:781-999-4286E_mail:info@targetmol.comAddress:36 Washington Street,Wellesley Hills,MA 02481