

Echistatin

Chemical Properties

CAS No. : 154303-05-6
Formula: C₂₁₇H₃₄₁N₇₁O₇₄S₉
Molecular Weight: 5417.1
Appearance: no data available
Storage: keep away from moisture
Powder: -20°C for 3 years | In solvent: -80°C for 1 year

EEESGPCRNCKFLKEGTICKRARGDDDDOYCNKGTCDPCRNPKGPAT
(Disulfide bridge Cys₉₂-Cys₁₁₁, Cys₉₂-Cys₉₂, Cys₉₂-Cys₉₂, Cys₉₂-Cys₉₂)

Biological Description

Description	Potent irreversible α V β 3 integrin antagonist (K_i = 0.27 nM). Disrupts attachment of osteoclasts to bone and inhibits bone reabsorption (IC_{50} = 0.1 nM). Prevents ADP-induced platelet aggregation via inhibition of glycoprotein IIb/IIIa (GpIIb/IIIa, α IIb β 3) receptors (IC_{50} = 30 nM) in vitro.
-------------	---

Solubility Information

Solubility	H ₂ O: 1 mg/mL, (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	0.1846 mL	0.923 mL	1.846 mL
5 mM	0.0369 mL	0.1846 mL	0.3692 mL
10 mM	0.0185 mL	0.0923 mL	0.1846 mL
50 mM	0.0037 mL	0.0185 mL	0.0369 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Reference

Musial et al (1990) Inhibition of platelet adhesion to surfaces of extracorporeal circuits by disintegrins RGD-containing peptides from viper venoms. Circulation 82 261 PMID:

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 36 Washington Street, Wellesley Hills, MA 02481