

A-674563 2HCl(552325-73-2(fb-2hcl))

## Chemical Properties

CAS No.:	T6139L
Formula:	C22H24Cl2N4O
Molecular Weight:	431.36
Appearance:	N/A
Storage:	0-4°C for short term (days to weeks), or -20°C for long term (months).

## Biological Description

Description	A-674563 is an Akt1 inhibitor with Ki of 11 nM, modest potent to PKA and >30-fold selective for Akt1 over PKC.
In vitro	A-674563 reduces phosphorylation of Akt downstream targets in cells and slows proliferation of tumor cells in vitro (EC50: 0.4 µM) [1]. A563 (0-10 µM) markedly decreases GSK3 and MDM2 phosphorylation in STS cells. A563 shows inhibitory effect on all STS cell lines, with IC50 values at 48 hours ranging from 0.22 µM to 0.35 µM. A563 induces G2 cell cycle arrest and apoptosis in STS cells. A563 (1 µM/12 hr) upregulates the expression of GADD45A independent of p53[2]. A-674563 (10-1000 nM) is anti-proliferative and cytotoxic in cultured human melanoma cells, induces melanoma cell apoptotic death, inhibited by caspase inhibitors, and inhibits melanoma cells via Akt-dependent and -independent mechanisms[3]. A-674563 is cytotoxic and anti-proliferative when added to U937 and Aml progenitor cells activate caspase-3/9 and apoptosis in U937 and Aml progenitor cells and manipulate other signalings in Aml cells whiling blocking Akt.
In vivo	In the PC-3 prostate cancer xenograft model, A-674563 (40 mg/kg/d, p.o.) has no marked monotherapy activity, but the efficacy of the combination therapy (A-674563+paclitaxel) is significantly improved. In an oral glucose tolerance test, A-674563 (20, 100 mg/kg) can increase plasma insulin[1]. A563 (20 mg/kg/bid; p.o.) exhibits slow tumor growth and a significant difference in tumor volume without significant weight loss of mice. A563-treated tumors express increased levels of GADD45α and decreased levels of PCNA (a nuclear marker for proliferation). Additionally, TUNEL assay staining levels (marker for apoptosis) increase in the A563-treated specimens[2]. A-674563 (25, 100 mg/kg, lavage daily) effectively inhibits A375 xenograft growth in mice[3]. A-674563 (15, 40 mg/kg) injection inhibits U937 xenograft in vivo growth and improves mice survival.
Cell Research	The cells on 96-well plates are gently washed with 200 µL of PBS. Alamar Blue reagent is diluted 1:10 in normal growth media. The diluted Alamar Blue reagent (100 µL) is added to each well on the 96-well plates and incubated until the reaction is complete. The analysis is done using a fmax Fluorescence Microplate Reader, set at the excitation wavelength of 544 nm and the emission wavelength of 595 nm. Data are analyzed using SOFTmax PRO software.

Animal Research	A-674563 is formulated in 5% dextrose. Immunocompromised male SCID mice are at 6 to 8 weeks of age. The 1×10 <sup>6</sup> 3T3-Akt1 or 2×10 <sup>6</sup> MiaPaCa-2 and PC-3 cells in 50% Matrigel are inoculated s.c. into the flank. For early treatment studies, mice are randomly assigned to treatment groups and therapy is initiated the day after inoculation. Ten animals are assigned to each group, including controls. For established tumor studies, tumors are allowed to reach a designated size and mice are assigned to treatment groups of equal tumor size (n=10 mice per group). Tumor size is evaluated by twice-weekly measurements with digital calipers. Tumor volume is estimated using the formula: V=L×W <sup>2</sup> /2. A-443654 is given s.c. in a vehicle of 0.2% HPMC. A-674563 is given orally in a vehicle of 5% dextrose. Gemcitabine and paclitaxel are added to the assay.
-----------------	---

## Solubility Information

Solubility	< 1 mg/ml refers to the product slightly soluble or insoluble
------------	---

### Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.318 mL	11.591 mL	23.182 mL
5 mM	0.464 mL	2.318 mL	4.636 mL
10 mM	0.232 mL	1.159 mL	2.318 mL
50 mM	0.046 mL	0.232 mL	0.464 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. The storage conditions and period of the stock solution: - 80 °C for 6 months; - 20 °C for 1 month. Please use it as soon as possible.

### Reference

1. Luo Y, Shoemaker A R, Liu X, et al. Potent and selective inhibitors of Akt kinases slow the progress of tumors in vivo.[J]. Molecular Cancer Therapeutics, 2005, 4(6):977.
2. Zhu Q S, Ren W, Korchin B, et al. Soft Tissue Sarcoma Are Highly Sensitive to AKT Blockade: A Role for p53 Independent Up-regulation of GADD45α[J]. Cancer Research, 2008, 68(8):2895-2903.
3. Zou Y, Zhu Y R, Fan G, et al. Pre-clinical assessment of A-674563 as an anti-melanoma agent[J]. Biochemical & Biophysical Research Communications, 2016, 477(1):1-8.

### Inhibitors · Natural Compounds · Compound Libraries

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use.

Tel:781-999-4286

E-mail:info@targetmol.com

Address:36 Washington Street,Wellesley Hills,MA 02481